

Next-to-next-to-leading order post-Newtonian linear-in-spin binary Hamiltonians

Johannes Hartung^{*1}, Jan Steinhoff^{1,2}, and Gerhard Schäfer¹

¹ Theoretisch–Physikalisches Institut,
Friedrich–Schiller–Universität,

Max–Wien–Platz 1, 07743 Jena, Germany, EU

² Centro Multidisciplinar de Astrofísica (CENTRA), Departamento de Física,
Instituto Superior Técnico (IST), Universidade Técnica de Lisboa,
Avenida Rovisco Pais 1, 1049-001 Lisboa, Portugal, EU

February 28, 2013

Abstract

The next-to-next-to-leading order post-Newtonian spin-orbit and spin(1)-spin(2) Hamiltonians for binary compact objects in general relativity are derived. The Arnowitt-Deser-Misner canonical formalism and its generalization to spinning compact objects in general relativity are presented and a fully reduced matter-only Hamiltonian is obtained. Several simplifications using integrations by parts are discussed. Approximate solutions to the constraints and evolution equations of motion are provided. Technical details of the integration procedures are given including an analysis of the short-range behavior of the integrands around the sources. The Hamiltonian of a test-spin moving in a stationary Kerr spacetime is obtained by rather simple approach and used to check parts of the mentioned results. Kinematical consistency checks by using the global (post-Newtonian approximate) Poincaré algebra are applied. Along the way a self-contained overview for the computation of the 3PN ADM point-mass Hamiltonian is provided, too.

PACS numbers: 04.25.Nx, 04.20.Fy, 04.25.-g, 97.80.-d, 45.50.Jf

Keywords: post-Newtonian approximation; canonical formalism; approximation methods; equations of motion; binary stars

1 Introduction

Since Einstein discovered the theory of general relativity [1–5], many attempts to solve the field equations were undertaken. Yet only a few analytical solutions to the full field equations are known at the time being, mostly for highly symmetric matter or field configurations [6, 7]. The most famous solutions of the field equations are the Schwarzschild [8] and the Kerr [9] ones. Even so, an analytic solution for binary systems of black holes (or even neutron stars) is missing and unlikely to be found in the future. Nevertheless, such binaries are very interesting. In particular they constitute the most promising and strongest sources for gravitational waves, one of the most fascinating predictions of the theory of general relativity [10, 11].

To observe gravitational waves one needs very sensitive detectors because of the tiny cross section of the waves with matter. There exist several ground-based detector projects like e.g. Geo600, VIRGO, and LIGO [12–14] for this purpose. Their sensitivity increased during the last years due to continuous upgrades and they probably will detect gravitational waves directly within the next few years. The mentioned large-scale detectors were started to be built after gravitational waves had been observed indirectly by measuring

^{*}Corresponding author E-mail: johannes.hartung@uni-jena.de, Phone: +49 3641 947 106, Fax: +49 3641 947 102

the change of the radial orbital period e.g. for the binary pulsar PSR 1913+16 by Hulse and Taylor [15] in 1978 (Nobel prize 1993). Further precise observations of orbital period decays are the *double* pulsar PSR J0737–3039, [16, 17], and the white dwarf binary J0651+2844 [18]. Their theoretical prediction is based on the famous quadrupole formula (see e.g. [19, Eq. (4.12)]) which gives an expression for the orbital averaged energy loss of the whole system due to gravitational radiation and can thus be translated into a decay of the orbital period.

If gravitational wave signals are detected in the near future, a great effort in data analysis will be necessary to extract them from the noisy raw data. Necessary ingredients to achieve this goal on the way towards gravitational wave astronomy are predictions on which kind of signals can be expected, e.g. from binary sources. The theoretical form of the signals – which should incorporate all known physical effects to some specific order of magnitude – is called template and relies on a solution of the field equations and therefore on the physical parameters of the system. Unfortunately, after around one hundred years of research there are no dynamical analytical solutions of the field equations for an n -body system ($n \geq 2$) known. There are two possibilities to circumvent this problem. The first one is to rely on numerical simulations. The second possibility is to rely on approximation methods to extract solutions for these kind of problems from the field equations. These include in particular the post-Minkowskian approximation, the post-Newtonian (PN) approximation, extreme mass ratios (including testmass case), and the effective-one-body approach.

One of the most successful approximation methods is the post-Newtonian approximation, a slow motion and wide separation approximation. It is used to treat the finite propagation velocity of the gravitational field approximately as an instantaneous effect and therefore “freeze” its dynamics. Thus the field degrees of freedom are eliminated in this approximation scheme. Afterwards one is left with ordinary differential equations for positions, momenta, and spins of the objects in the system only. It is convenient to encode these equations of motion into a Lagrangian or a Hamiltonian.

To get a more quantitative understanding of the post-Newtonian approximation, consider a gravitationally bound binary system in the Newtonian limit. In this case one can relate the Newtonian kinetic energy and the Newtonian gravitational potential through the Virial theorem, namely

$$\frac{v^2}{c^2} \sim \frac{GM}{c^2 r} \sim \frac{r_s}{r}, \quad (1)$$

with v denoting a typical orbital velocity in the system, c meaning the speed of light, G Newton’s gravitational constant, M the total mass of the system, r a typical distance between the gravitating objects, and r_s being the Schwarzschild radius of the total system. Every order in v^2/c^2 is denoted as one relative post-Newtonian order. The post-Newtonian approximation was used to obtain matter equations of motion for some special systems already shortly after general relativity had been developed, see e.g. the 1PN binary Lagrangian in [20] and the 1PN equations of motion in e.g. [21]. Furthermore previous results for non-spinning objects obtained within the formalism used in the present article, namely the canonical formalism of Arnowitt, Deser, and Misner (ADM) [22], are the 2PN [23–25], 2.5PN [19, 26], 3PN [27–31], and 3.5PN [32, 33] Hamiltonians. For various other (non-canonical) derivations of post-Newtonian results for non-rotating objects see [34–42] and references therein.

Regarding the spin corrections to the post-Newtonian approximation, the leading order can be found in [43–46]. Interestingly the leading-order equations of motion were already obtained earlier within the (more general) post-Minkowskian approximation [47, 48]. The post-Newtonian next-to-leading order in spin was only tackled more recently [49–58]. The post-Newtonian next-to-next-to-leading order spin-orbit and spin(1)-spin(2) Hamiltonians are the subject of the present paper. For very rapidly rotating objects they can be comparable in strength to 3.5PN and 4PN corrections, respectively. Half a post-Newtonian order above them the first (leading-order) spin-dependent *radiative* Hamiltonians appear. The spin-orbit and spin(1)-spin(2) ones were already obtained recently [59]. At the 3.5PN level one should further include all Hamiltonians cubic in the spins derived in [60, 61]. Notice that these cubic Hamiltonians are only known for binary black holes so far, whereas all other mentioned Hamiltonians (including the ones derived in the present paper) are valid for general compact objects [or have been generalized to this case, see [62–65] for the spin(1)-spin(1) level]. However, the Hamiltonians in the present paper are only obtained for binary systems (but many results for three and more objects already exist [66–70]). Besides spin effects, tidal

contributions to the post-Newtonian approximation become very important for general compact objects like neutron stars [71–73], also see e.g. [74] for the extreme mass ratio case.

The present article provides a detailed exposition of how the next-to-next-to-leading order (NNLO) spin-orbit and spin(1)-spin(2) post-Newtonian Hamiltonians can be derived from an extension of the canonical formalism of Arnowitt, Deser, and Misner [22]. The mentioned extension of the ADM formalism refers to a generalization from (non-rotating) point-masses (PM) to rotating objects [75], see also [76–78]. Results for these Hamiltonians were already presented in this journal [79, 80]. Their technicalities are also discussed in the (german) PhD thesis of JH [81]. A corresponding Lagrangian potential for the NNLO spin(1)-spin(2) interaction was derived simultaneously by M. Levi in [82] via an effective field theory (EFT) approach. A comparisons between EFT and ADM results at NNLO spin(1)-spin(2) was not yet undertaken, as it is not straightforward (see [83, 84] for a discussion at NLO). However, in [85, 86] the equations of motion in harmonic gauge were calculated at NNLO spin-orbit level and agreement with our Hamiltonian was found.

For unbound systems one can in general not relate the velocities of the objects to the strength of the gravitational coupling, i.e. the basic relation (1) of the post-Newtonian approximation is not applicable. For these kinds of systems (e.g. scattering of black holes) an useful approximation is the so called post-Minkowskian approximation, which is an expansion in powers of the gravitational coupling constant G only and thus also appropriate for very high velocities and weak fields. The first post-Minkowskian approximation for non-rotating objects was used to derive the Hamiltonian in [87] within the ADM formalism. In principle the expressions given in [26] can be used to derive the ADM Hamiltonian in the second post-Minkowskian approximation if there is no incoming radiation. But the integrals have not been given in closed form yet and since we are only interested in gravitationally bound systems, we retreat to the post-Newtonian approximation.

The Hamiltonians derived in this article are not the end of the journey. In order to extract useful information (i.e. the parameters of a binary) from the gravitational waves, one needs the solution of the post-Newtonian equations of motion for the binary.¹ For known orbital parameterizations one can further calculate the far-zone radiation field (see e.g [98–100] for general formalism of treating radiation in general relativity, [101] for higher order radiation losses in point-mass binaries, [102–104] for spin effects on the radiation, [105] for spin-dependent tail effects, and [106, 107] for multipole moments including spin up to 2.5PN). In case of eccentric orbits the radiation consists of several modes which may be extracted by a mode decomposition (see e.g. [108–110] for decomposition of the radiation field in tensor spherical harmonics, and computing and solving ordinary differential equations for mean motion and eccentricity in a binary without spin; see also [111–117] for higher order mode decomposition of multipole moments also for point-mass binaries only). Further for known parameterizations one is also able to calculate the loss of energy and angular momentum during the inspiral process, see e.g. [114, 118, 119] for time evolution effects on the template banks and [59, 120] for the near-zone luminosity. This is necessary to construct the mentioned template banks to extract the physical parameters from a noisy signal via a matched filtering procedure. There the “fitting factor” is a very sensitive indicator for the performance of the template bank vis-à-vis the real signal. An introduction to matched filtering can be found in [121, 122]. There exists a plethora of articles referring to circular inspiral without spin, e.g. [123–126].

Though the effects considered in the present article are very small, they still probably have a serious impact on future template banks. The reason for this is that even tiny contributions to the binary’s interaction accumulate during the long inspiral phase (where the post-Newtonian approximation is still valid) and thus may become observable for potentially planned space based detectors in the future. During the very late inspiral phase these effects will become more important, but the post-Newtonian approximation will break down due to the highly nonlinear behavior of the dynamics and high velocities ($v/c \gtrsim 1/3$). To overcome this problem it is most convenient to extrapolate to this nonlinear regime by resumming the post-Newtonian

¹The following literature and in particular [88] gives a complete overview over the research area of parameterization. See e.g. [89] for a point-mass 1PN parameterization, [90] for a point-mass 2PN parameterization, [91] for a quasi-Keplerian 3PN point-mass parameterization, [92, 93] for point-mass parameterizations under leading order spin-orbit coupling, [94] for using a 3PN point-mass parameterization including radiative dynamics for the phasing of gravitational waves, [95] for a post-equal-mass parameterization of a binary at 3PN point-mass level under leading order spin-orbit coupling, and finally [96] for a parameterization up to 2.5PN with orbital angular momentum aligned spins. [97] will incorporate the linear-in-spin Hamiltonians given in the present article into the orbital elements for orbital momentum aligned spins.

series. Such a resummation was successfully implemented into the effective-one-body approach, which analytically provides complete gravitational waveforms for binary inspiral that are in good agreement with numerical relativity. This succeeded so far for point-masses [127, 128] and for non-precessing spins [129] by calibrations to full numerical simulations, but more work is needed for precessing spins [130, 131]. Here the Hamiltonians derived in the present paper should be very useful, and the spin-orbit one was indeed already incorporated into the effective-one-body approach [132, 133]. See also [134] for a very complete overview of the literature on the effective-one-body approach. Alternative ways of resumming the post-Newtonian series by Padé approximants are possible, which is most interesting for certain gauge invariant quantities [135, 136]. Within the overlap region of post-Newtonian approximation and numerical relativity in which the gravitational field is not too strong and the number of orbits can be handled by numerical simulations the results of both approaches can be compared. The mentioned resummation methods can make these approximate results competitive to numerical relativity² also in the late inspiral phase [127, 128]. Last but not least, a powerful interface between self-force calculations and the ADM Hamiltonians derives from the redshift observable and the first law of binary dynamics [150], which was extended to include spin recently [151].

For all computations we used xTENSOR [152], a free package for MATHEMATICA [153], especially because of its fast index canonicalizer based on the package xPERM [154]. We also used the package xPERT [155], which is part of xTENSOR, for performing the perturbative part of our calculations. Furthermore we wrote several MATHEMATICA packages ourselves for the various steps of the computation including evaluating integrals.

The article is organized as follows. Sect. 2 shows how to split the spacetime in a $(d+1)$ manner, derives the Hamilton constraint and the momentum constraint from the Einstein-Hilbert action, and shows the definition of the ADM Hamiltonian. In Sect. 3 the constraint equations are expanded using a post-Newtonian power counting scheme and several integrations by parts simplify subsequent integrations. Afterwards a transition from the ADM Hamiltonian to a Routhian via a Legendre transform is performed in Sect. 4. The integrands are composed of various fields and their sources. These are explained in detail in Sect. 5. From the sources one can obtain the solution of the constraint equations and wave equations in Sect. 6 and is ready to integrate. Due to its importance a detailed explanation of the ultraviolet analysis is also provided there. In Sect. 7 the resulting next-to-next-to-leading order Hamiltonians are given. The kinematic consistency check of the Hamiltonians, namely the global post-Newtonian approximate Poincaré algebra, is discussed in Sect. 8. Also the center-of-mass vectors are uniquely determined from an ansatz in the same section. Another check is provided in Sect. 9. There the Hamiltonians from Sect. 7 are compared with the Hamiltonian of a test-spin in a stationary exterior gravitational field. After that in Sect. 10 the conclusions are presented and in the Appendices more details on some of the calculation procedures used in the former sections are provided.

The spacetime has d spatial dimensions, 1 time dimension, and metric signature $d - 1$. In this article a restriction to $d = 3$ is explicitly written, if this is not the case d is always generic. All calculations at the level of the field equations are performed in arbitrary dimensions, because of the necessary ultraviolet(UV) analysis concerning the short-range decay of fields around the sources. Vectors are written in boldface and their components are denoted by Latin indices from the middle of the alphabet. The scalar product between two vectors $\mathbf{a}$ and $\mathbf{b}$ is denoted by $(\mathbf{a} \mathbf{b}) \equiv (\mathbf{a} \cdot \mathbf{b})$. Our units are such that $c = 1$. There is no special convention for Newton's gravitational constant $G^{(d)}$. (Notice that $G^{(d)}$ is the d -dimensional coupling strength of the gravitational field. It has the same numerical value as G in $d = 3$ dimensions, but different units.) In the results and the expressions for the sources, $\hat{\mathbf{p}}_a$ denotes the canonical linear momentum of the a th object, $\hat{\mathbf{z}}_a$ the canonical conjugate position of the object, m_a the mass of the object, $\hat{\mathbf{S}}_a$ and $\hat{S}_{a(i)(j)}$ the spin vector and the spin tensor of the object, $r_{ab} = |\hat{\mathbf{z}}_a - \hat{\mathbf{z}}_b|$ the relative distance between two objects, and $\mathbf{n}_{ab} = (\hat{\mathbf{z}}_a - \hat{\mathbf{z}}_b)/r_{ab}$ the direction vector pointing from object b to object a . Also important are the distance between source a and field point, $r_a = |\mathbf{x} - \hat{\mathbf{z}}_a|$, the unit vector pointing from source a to the field point $\mathbf{n}_a = (\mathbf{x} - \hat{\mathbf{z}}_a)/r_a$, and the circumference of the triangle of source a , source b and the field point $\mathbf{x}$, given by $s_{ab} = r_a + r_b + r_{ab}$. In the binary case the object labels a, b take only the values 1 and 2. The round brackets around the indices of the canonical spin tensor $\hat{S}_{a(i)(j)}$ indicate that its

²See e.g. [137–139] for reviews, [140] for the first simulation of binary neutron stars, and e.g. [141–143] for very recent studies about them. The first simulations of coalescences of binary black holes were performed in [144, 145]. Furthermore see e.g. [146–148] for numerical simulations of a system of more than two black holes and the recent publication [149] about its chaotic behavior.

components are given in a local Lorentz basis, which is essential for the canonical formalism, see [75, 76].

2 ADM Canonical Formalism

In the following we introduce the ADM canonical formalism [22, 156, 157]. We pay special attention to the dependence on the spatial dimensions d , as dimensional regularization is important for the consistency of the post-Newtonian approximation when point-like sources are utilized [31]. The ADM formalism is extended from non-spinning to spinning point-like sources valid to linear order in spin here, see [75, 76] for the case $d = 3$.

2.1 Setting the Canonical Formalism

Canonical methods usually require one to single out a time coordinate, thus splitting spacetime into space and time. The geometrically favored way for such a splitting gives rise to the line element

$$ds^2 = -N^2 dt^2 + \gamma_{ij}(N^i dt + dx^i)(N^j dt + dx^j), \quad (2)$$

which corresponds to a $(d + 1)$ -dimensional metric tensor field given by

$$g_{\mu\nu} = \begin{pmatrix} -N^2 + N_i N^i & N_i \\ N_j & \gamma_{ij} \end{pmatrix}, \quad g^{\mu\nu} = \begin{pmatrix} -1/N^2 & N^i/N \\ N^j/N & \gamma^{ij} - \frac{N^i N^j}{N^2} \end{pmatrix}, \quad (3)$$

where N is the lapse function, N^i the shift vector, $N_i = \gamma_{ij} N^j$, and γ_{ij} the metric of the spatial slices, γ^{ij} being its inverse [158–160]. Notice that rewriting the metric tensor field in terms of lapse, shift, and the d -dimensional metric γ_{ij} corresponds only to another representation of the metric, since together they have the same number of degrees of freedom. On the one hand a symmetric rank two tensor field like $g_{\mu\nu}$ in $d + 1$ dimensions has $(d + 1)(d + 2)/2$ independent entries and on the other hand, a symmetric rank two tensor in d dimensions like γ_{ij} has $d(d + 1)/2$ degrees of freedom, the vector N^i has d independent entries, and the scalar N represents one degree of freedom. Obviously the degrees of freedom match.

The action of the gravitational field is given by the usual Einstein–Hilbert action, namely

$$W_{\text{field}} = \int d^{d+1}x \mathcal{L}_{\text{field}}, \quad \mathcal{L}_{\text{field}} = \frac{1}{16\pi G^{(d)}} \sqrt{-g}^{(d+1)} \mathbf{R}, \quad (4)$$

where $g = \det(g_{\mu\nu})$ and $^{(d+1)}\mathbf{R}$ is the $(d + 1)$ -dimensional Ricci scalar, which can be split in a $(d + 1)$ manner resulting in

$$\mathcal{L}_{\text{field}} = \frac{1}{16\pi G^{(d)}} N \sqrt{\gamma} \left[^{(d)}\mathbf{R} + K_{ij} K^{ij} - (\gamma_{ij} K^{ij})^2 \right] + (\text{td}). \quad (5)$$

Here $^{(d)}\mathbf{R}$ is the d -dimensional spatial Ricci scalar, K_{ij} is the extrinsic curvature, $\gamma = \det(\gamma_{ij})$, and (td) denotes a total divergence which we ignore for now (it will be discussed in Sect. 2.4). We define

$$\pi^{ij} = 16\pi G^{(d)} \frac{\partial \mathcal{L}_{\text{field}}}{\partial \gamma_{ij,0}} = \sqrt{\gamma} (\gamma^{ij} \gamma^{kl} - \gamma^{ik} \gamma^{jl}) K_{kl}, \quad (6)$$

where $2N K_{ij} = -\gamma_{ij,0} + 2N_{(i;j)}$ was used (which deviates from the convention for K_{ij} used in [159]), a comma denotes a partial derivative, and a semicolon denotes the d -dimensional covariant derivative. The inversion reads

$$K_{ij} = \frac{1}{\sqrt{\gamma}} \left(\frac{1}{d-1} \gamma_{ij} \gamma_{kl} - \gamma_{ik} \gamma_{jl} \right) \pi^{kl}, \quad (7)$$

and the dimension finally enters the calculation explicitly. In order to put the field equations in the form of Hamilton’s canonical equations, we perform a Legendre transform,

$$\mathcal{L}_{\text{field}} = \frac{1}{16\pi G^{(d)}} \pi^{ij} \gamma_{ij,0} - N \mathcal{H}^{\text{field}} + N^i \mathcal{H}_i^{\text{field}} + (\text{td}), \quad (8)$$

where

$$\mathcal{H}^{\text{field}} = -\frac{1}{16\pi\sqrt{\gamma}G^{(d)}} \left[\gamma^{(d)}\mathbf{R} - \gamma_{ij}\gamma_{kl}\pi^{ik}\pi^{jl} + \frac{1}{d-1} (\gamma_{ij}\pi^{ij})^2 \right], \quad (9a)$$

$$\mathcal{H}_i^{\text{field}} = \frac{1}{8\pi G^{(d)}} \gamma_{ij}\pi^{jk}_{;k}. \quad (9b)$$

Variation of the action with respect to γ_{ij} and π^{ij} leads to $d(d+1)$ first order evolution equations for these variables. The d -metric γ_{ij} and the field momentum $\frac{1}{16\pi G^{(d)}}\pi^{ij}$ are canonically conjugate variables in the vacuum case. However, the action does not contain time derivatives of N and N^i , so a variation with respect to these variables leads to constraint-type equations (i.e. containing no time derivatives). Variation of N gives only one equation called the Hamilton constraint, while variation of N^i gives d equations called momentum constraint. We have arrived at a canonical formalism without any restriction on the coordinates, or gauge, but in the presence of constraints. In the following sections we will eliminate these constraints by simultaneously fixing the gauge.

In order to couple the ADM formalism to point-like objects possessing masses m_a and $(d+1)$ -dimensional spin tensors $S_{a\mu\nu} = -S_{a\nu\mu}$, it is best to start with a matter action of the form

$$W_{\text{matter}} = \sum_a \int d\tau_a \left[p_{a\mu} u_a^\mu + \frac{1}{2} S_{a\mu\nu} \Omega_a^{\mu\nu} - \lambda_a (g^{\mu\nu} p_{a\mu} p_{a\nu} + m_a^2) \right], \quad (10)$$

where τ_a is a worldline parameter, $p_{a\mu}$ the $(d+1)$ -momentum, $u_a^\mu = dz_a^\mu/d\tau_a$, z_a^μ the position, $\Omega_a^{\mu\nu} = -\Omega_a^{\nu\mu}$ the angular velocity tensor, and λ_a a Lagrange multiplier belonging to the mass-shell constraint $g^{\mu\nu} p_{a\mu} p_{a\nu} + m_a^2 = 0$ (all for the a -th object), see [75, 76] for details. Notice that the angular velocity tensor is built from a Lorentz matrix encoding the orientation of a body-fixed frame and its covariant derivative. This means one has to handle a derivative coupling of the metric because of Christoffel symbols or Ricci rotation coefficients appearing in the covariant derivative of the Lorentz matrix. Further constraints must be fulfilled for the mentioned Lorentz matrix and the spin, the latter reads $S_a^{\mu\nu} p_{a\mu} = 0$. It is important that this action is valid to linear order in spin and is now considered for a generic spatial dimension d . The equations of motion following from this matter action are the well-known Mathisson-Papapetrou equations [161, 162], see also [163, 164],

$$\frac{DS_a^{\mu\nu}}{d\tau_a} = 2p_a^{[\mu} u_a^{\nu]}, \quad \frac{Dp_{a\mu}}{d\tau_a} = -\frac{1}{2} {}^{(d+1)}\mathbf{R}_{\mu\rho\beta\alpha} u_a^\rho S_a^{\beta\alpha}, \quad (11)$$

where ${}^{(d+1)}\mathbf{R}_{\mu\rho\beta\alpha}$ is the $(d+1)$ -dimensional Riemann tensor, and the source of the gravitational field equations is given by Tulczyjew's singular energy-momentum tensor density [163]

$$\sqrt{-g}T^{\mu\nu} = \sum_a \int d\tau_a \left[u_a^{(\mu} p_a^{\nu)} \delta_{(d+1)a} - \left(S_a^{\alpha(\mu} u_a^{\nu)} \delta_{(d+1)a} \right)_{\parallel\alpha} \right], \quad (12)$$

where $\delta_{(d+1)a} = \delta(x^\mu - z_a^\mu)$. For a review of spin in relativity see e.g. [165, 166]. Further details on a corresponding action principle can be found in [167–171].

Next the matter constraints are eliminated from the action with suitable gauge choices, e.g. $\tau_a = t$ where t is the coordinate time and the matter variables are transformed to (reduced) canonical variables denoted by a hat, e.g. $\hat{z}_a^i$ or $\hat{p}_{ai}$. This is completely analogous to [75, 76] and will therefore not be repeated here. But the following facts are important for the present article. The spatial dimension d is not entering the just mentioned calculations explicitly, whereas it appears in the gravitational part of the action. Further, the matter action becomes linear in lapse and shift, so the gravitational constraints following from their variation now contain matter source terms $\mathcal{H}^{\text{matter}}$ and $\mathcal{H}_i^{\text{matter}}$,

$$\mathcal{H}^{\text{field}} + \mathcal{H}^{\text{matter}} = 0, \quad \mathcal{H}_i^{\text{field}} + \mathcal{H}_i^{\text{matter}} = 0. \quad (13)$$

Finally, due to the coupling of the spin to derivatives of the metric, time derivatives of the gravitational field are present in the matter part. This necessitates a matter contribution π_{matter}^{ij} to the *canonical* field momentum, which now reads

$$\hat{\pi}^{ij} = \pi^{ij} + \pi_{\text{matter}}^{ij}. \quad (14)$$

Explicit expressions for $\mathcal{H}^{\text{matter}}$, $\mathcal{H}_i^{\text{matter}}$, and π_{matter}^{ij} are provided in Sect. 5 and have the same form as for $d = 3$ in [75, 76].

The next goal is the elimination of the field constraints within the so called ADM transverse-traceless gauge (ADMTT). The corresponding gauge conditions read

$$\gamma_{ij,j} - \frac{1}{d}\gamma_{jj,i} = 0, \quad (15a)$$

$$\hat{\pi}^{ii} = 0. \quad (15b)$$

We proceed to work out field decompositions suitable for an elimination of the field constraints.

2.2 Metric Decomposition

The differential gauge condition (15a) for the metric is solved by

$$\gamma_{ij} = \Psi\delta_{ij} + h_{ij}^{\text{TT}}, \quad (16)$$

which can be seen from a transverse-traceless (TT) decomposition of γ_{ij} . The first part is the conformally flat part of the metric and the last part can be interpreted in the far-zone as the radiation field, which is transverse-traceless, i.e. $h_{ii}^{\text{TT}} = 0 = h_{ij,j}^{\text{TT}}$. Due to the requirement of maximal simple curvature density expression [172], one can determine the conformal part of the metric in a form appropriate for a post-Newtonian expansion. Consider the static (i.e. momentum independent) part of the Hamilton constraint using (9a) and (13) setting $\pi^{ij} = 0$,

$$\sqrt{\gamma}^{(d)}\text{R} = 16\pi G^{(d)}\mathcal{H}^{\text{matter}}. \quad (17)$$

If one inserts the truncation $\gamma_{ij} = \Psi\delta_{ij}$ with the ansatz $\Psi = \psi^\beta$ into the static Hamilton constraint, one obtains

$$-\frac{1}{4}\beta(d-1)\psi^{-2+\frac{1}{2}\beta(d-2)}(4\psi\Delta\psi + (-4 + \beta(d-2))(\psi_{,i})^2) = 16\pi G^{(d)}\mathcal{H}^{\text{matter}}, \quad (18)$$

where $\Delta = \partial_i\partial_i$ and ∂_i denotes a partial coordinate derivative. Demanding that the nonlinear gradient term $(\psi_{,i})^2$ disappears (yielding a Poisson-type equation) or the ψ in front has a vanishing exponent, both leads to

$$\beta = \frac{4}{d-2}, \quad (19)$$

and thus gives the most simple expression for the curvature density $\sqrt{\gamma}^{(d)}\text{R}$. Using this solution, the static Hamilton constraint reduces to

$$-\frac{4(d-1)}{d-2}\psi\Delta\psi = 16\pi G^{(d)}\mathcal{H}^{\text{matter}}. \quad (20)$$

Now one can further set $\psi = 1 + \alpha\phi$ and demand that the static part of the Hamilton constraint linear in ϕ reduces to a Poisson-type equation without any d -dependence. This leads to

$$\alpha = \frac{d-2}{4(d-1)}, \quad (21)$$

and

$$-\left(1 + \frac{d-2}{4(d-1)}\phi\right)\Delta\phi = 16\pi G^{(d)}\mathcal{H}^{\text{matter}}. \quad (22)$$

Finally, the optimized metric decomposition reads

$$\gamma_{ij} = \left(1 + \frac{d-2}{4(d-1)}\phi\right)^{4/(d-2)}\delta_{ij} + h_{ij}^{\text{TT}}, \quad (23)$$

see, e.g., [31]. One can also borrow the conformal factor from the d -dimensional isotropic Schwarzschild metric given in, e.g., [173, Eq. (22)]. For convenience we introduce

$$\bar{\phi} = \frac{d-2}{4(d-1)}\phi, \quad (24)$$

in order to absorb certain dimension-dependent factors in subsequent calculations.

2.3 Momentum Decomposition

For convenience we introduce some differential-integral operators called $\mathcal{D}$ -operators, given by

$$\mathcal{D}_a^{ij} = \delta_{ij} + (a-1)\partial_i\partial_j\Delta^{-1} = \mathcal{D}_a^{ji}, \quad (25)$$

where Δ^{-1} is the inverse Laplacian with usual boundary conditions. These have the following nice properties

$$\frac{1}{n} \sum_n \mathcal{D}_{a_n}^{ij} = \mathcal{D}_{\frac{1}{n} \sum_n a_n}^{ij}, \quad (26a)$$

$$\mathcal{D}_a^{ik} \mathcal{D}_b^{kj} = \mathcal{D}_{ab}^{ij}, \quad (26b)$$

$$\mathcal{D}_{a-1}^{ij} = (\mathcal{D}_a^{ij})^{-1}, \quad (26c)$$

and further

$$\mathcal{D}_1^{ij} = \delta_{ij}, \quad (27a)$$

$$\delta_{ij} \mathcal{D}_a^{ij} = d + a - 1, \quad (27b)$$

$$\partial_i \mathcal{D}_a^{ij} = a \partial_j. \quad (27c)$$

Note that for $a = 0$ one gets the transverse projector

$$\perp^{ij} := \mathcal{D}_0^{ij}, \quad (28)$$

which has vanishing divergence and is not invertible. From the multiplication property of the $\mathcal{D}$ -operators the projection property $\perp^{ik} \perp^{kj} = \perp^{ij}$ follows immediately. Also a longitudinal projector can be constructed, namely

$$\parallel^{ij} := \mathcal{D}_1^{ij} - \mathcal{D}_0^{ij} = \delta_{ij} - \perp^{ij}. \quad (29)$$

Obviously the longitudinal projector also fulfills the projector condition $\parallel^{ik} \parallel^{kj} = \parallel^{ij}$. According to [76, 77] one can decompose the field momentum as

$$\pi^{ij} = \pi_{\text{TT}}^{ij} + \tilde{\pi}^{ij} + \check{\pi}^{ij}. \quad (30)$$

The different parts are given by

$$\pi_{\text{TT}}^{ij} = \left[\perp^{k(i} \perp^{j)\ell} - \frac{1}{d-1} \perp^{ij} \perp^{k\ell} \right] \pi^{k\ell} =: \delta_{ij}^{\text{TT} k\ell} \pi^{k\ell}, \quad (31a)$$

$$\tilde{\pi}^{ij} = \left[\parallel^{k(i} \delta_{j)\ell} + \delta_{k(i} \parallel^{j)\ell} - \parallel^{k(i} \parallel^{j)\ell} - \frac{1}{d-1} \perp^{ij} \parallel^{k\ell} \right] \pi^{k\ell} =: \delta_{ij}^{\text{LT} k\ell} \pi^{k\ell}, \quad (31b)$$

$$\check{\pi}^{ij} = \frac{1}{d-1} \perp^{ij} \delta_{k\ell} \pi^{k\ell} =: \delta_{ij}^{\text{Tr} k\ell} \pi^{k\ell}. \quad (31c)$$

Obviously π_{TT}^{ij} has no divergence and is trace free (it is transverse-traceless), $\tilde{\pi}^{ij}$ is trace free and its divergence contains the divergence of the whole field momentum, and $\check{\pi}^{ij}$ is divergence free and its trace

is the trace of the whole field momentum. This decomposition is essentially the usual transverse-traceless-decomposition of a symmetric rank two tensor field, but rearranged in a way more suitable for the present computations. For example, $\tilde{\pi}^{ij}$ is fixed by the gauge condition (15b) as

$$\tilde{\pi}^{ij} = -\frac{1}{d-1} \perp^{ij} \pi_{\text{matter}}^{kk}, \quad (32)$$

which follows from its definition and the trace of Eq. (14). Furthermore one can decompose $\tilde{\pi}^{ij}$ into two different vector potentials, $\tilde{\pi}^i$ and V^i . The decompositions read

$$\tilde{\pi}^{ij} = \tilde{\pi}_{,j}^i + \tilde{\pi}_{,i}^j - \frac{1}{d-1} \mathcal{D}_{d-1}^{ij} \tilde{\pi}_{,k}^k, \quad (33)$$

$$= V_{,j}^i + V_{,i}^j - \frac{2}{d} \delta_{ij} V_{,k}^k. \quad (34)$$

Of course $\tilde{\pi}^i$ and V^i are interrelated,

$$\tilde{\pi}^i = \mathcal{D}_{\frac{d-1}{2}}^{ij} V^j, \quad V^i = \mathcal{D}_{\frac{d}{2(d-1)}}^{ij} \tilde{\pi}^j, \quad (35)$$

and it holds

$$\tilde{\pi}^i = \Delta^{-1} \tilde{\pi}_{,j}^{ij}. \quad (36)$$

These two vector potentials have different advantages. From V^i it is possible to compute $\tilde{\pi}^{ij}$ without any integration. On the other hand $\tilde{\pi}^i$ has a simpler structure and can more easily be obtained from the momentum constraint using (36) [cf. (113), (114), and (115)]. The transverse-traceless parts of metric γ_{ij} , h_{ij}^{TT} and of the canonical field momentum $\hat{\pi}^{ij}$, $\hat{\pi}_{\text{TT}}^{ij}$ are denoted as transverse-traceless degrees of freedom or propagating field degrees of freedom in the following. The latter denotation will become clear in the next section.

2.4 ADM Hamiltonian

We are now principally able to solve the $d+1$ constraint equations (13) for the $d+1$ variables ϕ and $\tilde{\pi}^i$. Though this involves solving a system of nonlinear partial differential equations, it can be tackled analytically within the post-Newtonian approximation up to a certain order. Notice that $\tilde{\pi}^{ij}$ is fixed by (32) and that π_{TT}^{ij} can be replaced by

$$\pi_{\text{TT}}^{ij} = \hat{\pi}_{\text{TT}}^{ij} - \delta_{ij}^{\text{TT} k\ell} \pi_{\text{matter}}^{k\ell}. \quad (37)$$

The independent variables are thus the *reduced* canonical field variables h_{ij}^{TT} and $\hat{\pi}_{\text{TT}}^{ij}$ with Poisson brackets

$$\{h_{ij}^{\text{TT}}(\mathbf{x}), \hat{\pi}_{\text{TT}}^{k\ell}(\mathbf{x}')\} = 16\pi G^{(d)} \delta_{ij}^{\text{TT} k\ell} \delta(\mathbf{x} - \mathbf{x}'), \quad (38)$$

as well as reduced canonical matter variables which enter via the matter parts of the constraint equations (13) and are discussed in more detail in Sect. 5.

It was shown in [22, 156, 157] using different methods that the fully reduced Hamiltonian after gauge fixing is given by the ADM energy

$$E_{\text{ADM}} = \frac{1}{16\pi G^{(d)}} \oint d^{d-1} s_i [\gamma_{ij,j} - \gamma_{jj,i}], \quad (39)$$

where $\oint d^{d-1} s_i$ denotes an integral over the asymptotic boundary of a spatial hypersurfaces at fixed time. (The identical expression for the energy follows from the Landau-Lifshitz superpotential which is related with the well-known Landau-Lifshitz stress-energy-momentum pseudotensor of the gravitational field, see [174].) More precisely, the ADM energy E_{ADM} turns into the ADM Hamiltonian H_{ADM} if it is expressed in

terms of the mentioned reduced canonical variables. Inserting the metric decomposition (23), the surface integral can be transformed into a volume integral,

$$H_{\text{ADM}} = -\frac{1}{16\pi G^{(d)}} \int d^d x \Delta\phi, \quad (40)$$

where the asymptotic behavior of certain field components was used, see e.g. [157]. It was argued by Regge and Teitelboim [157] that one must modify the total divergence in (5) such that it leads to the surface integral expression for the ADM energy E_{ADM} , otherwise the variational principle is not well posed. Indeed, for the same reason one should add the (regularized) York–Gibbons–Hawking, or “trace K,” surface term [175–178] already to the Einstein–Hilbert action. The ADM energy then directly arises from a surface term contained in the complete action, see [179, 180] and also [159]. Notice that the Einstein equations can be followed from a variation of the action by disregarding all surface terms, but this is in general not allowed if the variation of the action has prescribed nontrivial behavior at the boundary.

The ADM Hamiltonian still depends on the reduced canonical field variables h_{ij}^{TT} and $\hat{\pi}_{\text{TT}}^{ij}$. In Sect. 4 also these remaining field variables will be eliminated through a Routhian approach to arrive at a conservative matter-only Hamiltonian.

3 Expansion of the Constraints and Integrations by Parts

As already stated in Sect. 1 we will use the post-Newtonian approximation throughout this article. Furthermore as mentioned above we utilize the spin-extended ADM formalism. So first of all we have to solve the constraint equations order by order. This requires to expand them in powers of the post-Newtonian approximation parameter (1) with the decompositions (23) and (30) (which are adapted to the ADMTT gauge) inserted. This is the task performed in the following subsection.

3.1 Order Counting

The field and source expansions starting at their leading order are given by

$$\phi = \phi_{(2)} + \phi_{(4)} + \phi_{(6)} + \phi_{(8)} + \phi_{(10)} + \dots, \quad (41a)$$

$$\tilde{\pi}^{ij} = \tilde{\pi}_{(3)}^{ij} + \tilde{\pi}_{(5)}^{ij} + \tilde{\pi}_{(7)}^{ij} + \dots, \quad (41b)$$

$$\mathcal{H}^{\text{matter}} = \mathcal{H}_{(2)}^{\text{matter}} + \mathcal{H}_{(4)}^{\text{matter}} + \mathcal{H}_{(6)}^{\text{matter}} + \mathcal{H}_{(8)}^{\text{matter}} + \mathcal{H}_{(10)}^{\text{matter}} + \dots, \quad (41c)$$

$$\mathcal{H}_i^{\text{matter}} = \mathcal{H}_{i(3)}^{\text{matter}} + \mathcal{H}_{i(5)}^{\text{matter}} + \mathcal{H}_{i(7)}^{\text{matter}} + \dots, \quad (41d)$$

where the subscript in round brackets denotes the c^{-1} order. A similar order counting is also valid for all derived fields, like vector potentials. At a later stage of the calculation we also need to expand h_{ij}^{TT} and $\hat{\pi}_{\text{TT}}^{ij}$,

$$h_{ij}^{\text{TT}} = h_{(4)ij}^{\text{TT}} + h_{(6)ij}^{\text{TT}} + \dots, \quad (41e)$$

$$\hat{\pi}_{\text{TT}}^{ij} = \hat{\pi}_{(5)\text{TT}}^{ij} + \dots \quad (41f)$$

At the mentioned stage we also need an order counting for the deviations between field momentum and canonical field momentum due to π_{matter}^{ij} and the tracelessness violation of the field momentum $\tilde{\pi}^{ij}$. As one can see in Sect. 5.3 π_{matter}^{ij} starts at $\mathcal{O}(c^{-9})$ and thus cannot contribute to the expansion of the Hamilton and momentum constraint later, while $\tilde{\pi}_{\text{matter}}^{ij}$ starts at $\mathcal{O}(c^{-5})$.

In anticipation of the source expressions in Sect. 5 we will introduce the matter variables used there to discuss the order counting of the field variables, because all fields depend on the source expression and thus the order counting of the matter variables. The mass m_a , canonical matter momentum $\hat{\mathbf{p}}_a$, and spin variables $\hat{\mathbf{S}}_a$ are *formally* counted as $m_a \sim \mathcal{O}(c^{-2})$, $\hat{\mathbf{p}}_a \sim \mathcal{O}(c^{-3})$, and $\hat{\mathbf{S}}_a \sim \mathcal{O}(c^{-3})$ for dimensional reasons only (remember that for *maximal* spins one would have $\hat{\mathbf{S}}_a \sim \mathcal{O}(c^{-4})$ instead, see e.g. [77, Appendix A]). This counting comes from the fact that after setting $c = G^{(d)} = 1$ we require all quantities to be in units of length. Let us introduce symbols with a bar below them being the quantities in SI units

and the other symbols the quantities in units of length, then it holds $m_a = \frac{G^{(d)}}{c^2} \underline{m}_a$ for the mass, $t = c\underline{t}$ for the time, $\hat{\mathbf{p}}_a = \frac{G^{(d)}}{c^3} \underline{\mathbf{p}}_a$ for the linear momentum, and similarly for the spin variables. Although we mentioned that $G^{(d)}$ has different units in $d \neq 3$ than in $d = 3$, we treat their c^{-1} order as in $d = 3$ for simplicity. So the order counting comes from the c powers inserted to reconstruct the SI units. It should be noted that these counting rules will in general not give correct *absolute* orders in c if the SI units of the final expression are not taken into account. However, *relative* orders are always meaningful, which is all that is relevant for perturbative expansions. Further notice that different counting rules are obtained if one assumes that all quantities are expressed in terms of mass units instead of length units when setting $c = G^{(d)} = 1$, which is also often used in the literature.

3.2 Hamilton Constraint

The Hamilton constraint

$$\frac{1}{16\pi\sqrt{\gamma}G^{(d)}} \left[\gamma^{(d)} \mathbf{R} - \gamma_{ij} \gamma_{kl} \pi^{ik} \pi^{jl} + \frac{1}{d-1} (\gamma_{ij} \pi^{ij})^2 \right] = \mathcal{H}^{\text{matter}}, \quad (42)$$

has to be expanded in a post-Newtonian manner. After inserting the metric and momentum decomposition, (23) and (30), and all field expansions (41) it decomposes into several Poisson equations for the post-Newtonian potential, namely

$$-\frac{1}{16\pi G^{(d)}} \Delta \phi_{(2)} = \mathcal{H}_{(2)}^{\text{matter}}, \quad (43a)$$

$$-\frac{1}{16\pi G^{(d)}} \Delta \phi_{(4)} = \mathcal{H}_{(4)}^{\text{matter}} - \bar{\phi}_{(2)} \mathcal{H}_{(2)}^{\text{matter}}, \quad (43b)$$

$$-\frac{1}{16\pi G^{(d)}} \Delta \phi_{(6)} = \mathcal{H}_{(6)}^{\text{matter}} - \bar{\phi}_{(2)} \mathcal{H}_{(4)}^{\text{matter}} + (-\bar{\phi}_{(4)} + \bar{\phi}_{(2)}^2) \mathcal{H}_{(2)}^{\text{matter}} - \frac{1}{16\pi G^{(d)}} \left(-(\tilde{\pi}_{(3)}^{ij})^2 + 4(\bar{\phi}_{(2)} h_{ij}^{\text{TT}})_{,ij} \right), \quad (43c)$$

$$\begin{aligned} -\frac{1}{16\pi G^{(d)}} \Delta \phi_{(8)} = & \mathcal{H}_{(8)}^{\text{matter}} - \bar{\phi}_{(2)} \mathcal{H}_{(6)}^{\text{matter}} + (-\bar{\phi}_{(4)} + \bar{\phi}_{(2)}^2) \mathcal{H}_{(4)}^{\text{matter}} + (-\bar{\phi}_{(6)} + 2\bar{\phi}_{(2)} \bar{\phi}_{(4)} \\ & - \bar{\phi}_{(2)}^3) \mathcal{H}_{(2)}^{\text{matter}} - \frac{1}{16\pi G^{(d)}} \left(\frac{3d-10}{d-2} \bar{\phi}_{(2)} (\tilde{\pi}_{(3)}^{ij})^2 - 2\tilde{\pi}_{(3)}^{ij} \tilde{\pi}_{(5)}^{ij} - 2\tilde{\pi}_{(3)}^{ij} \pi_{\text{TT}}^{ij} \right. \\ & + \frac{4}{d-2} h_{ij}^{\text{TT}} \bar{\phi}_{(2),i} \bar{\phi}_{(2),j} - \frac{1}{4} (h_{ij,k}^{\text{TT}})^2 - \frac{16}{d-2} (\bar{\phi}_{(2)} \bar{\phi}_{(2),j} h_{ij}^{\text{TT}})_{,i} + 4(\bar{\phi}_{(4)} h_{ij}^{\text{TT}})_{,ij} \\ & \left. + \frac{1}{2} \Delta (h_{ij}^{\text{TT}})^2 - \frac{1}{2} (h_{ik}^{\text{TT}} h_{jk}^{\text{TT}})_{,ij} \right), \end{aligned} \quad (43d)$$

$$\begin{aligned} -\frac{1}{16\pi G^{(d)}} \Delta \phi_{(10)} = & \mathcal{H}_{(10)}^{\text{matter}} - \bar{\phi}_{(2)} \mathcal{H}_{(8)}^{\text{matter}} + (-\bar{\phi}_{(4)} + \bar{\phi}_{(2)}^2) \mathcal{H}_{(6)}^{\text{matter}} + (-\bar{\phi}_{(6)} + 2\bar{\phi}_{(2)} \bar{\phi}_{(4)} \\ & - \bar{\phi}_{(2)}^3) \mathcal{H}_{(4)}^{\text{matter}} + \left(-\bar{\phi}_{(8)} + 2\bar{\phi}_{(2)} \bar{\phi}_{(6)} + \bar{\phi}_{(4)}^2 - 3\bar{\phi}_{(4)} \bar{\phi}_{(2)}^2 + \bar{\phi}_{(2)}^4 \right. \\ & \left. + \frac{d-4}{8(d-1)} (h_{ij}^{\text{TT}})^2 \right) \mathcal{H}_{(2)}^{\text{matter}} - \frac{1}{16\pi G^{(d)}} \left(\frac{3d-10}{d-2} \bar{\phi}_{(4)} (\tilde{\pi}_{(3)}^{ij})^2 - 2h_{ij}^{\text{TT}} \tilde{\pi}_{(3)}^{ik} \tilde{\pi}_{(3)}^{jk} \right. \\ & - (\tilde{\pi}_{(5)}^{ij})^2 - 2\tilde{\pi}_{(3)}^{ij} \tilde{\pi}_{(7)}^{ij} - (\pi_{\text{TT}}^{ij})^2 - 2\frac{(d-3)(3d-10)}{(d-2)^2} (\tilde{\pi}_{(3)}^{ij})^2 \bar{\phi}_{(2)}^2 \\ & + \frac{8}{d-2} h_{ij}^{\text{TT}} \bar{\phi}_{(2),i} \bar{\phi}_{(4),j} + 2\frac{3d-10}{d-2} \tilde{\pi}_{(3)}^{ij} \tilde{\pi}_{(5)}^{ij} \bar{\phi}_{(2)} + 2\frac{3d-10}{d-2} \tilde{\pi}_{(3)}^{ij} \pi_{\text{TT}}^{ij} \bar{\phi}_{(2)} \\ & \left. - 4\frac{d+2}{(d-2)^2} h_{ij}^{\text{TT}} \bar{\phi}_{(2),i} \bar{\phi}_{(2),j} \bar{\phi}_{(2)} - \frac{1}{2} h_{ij,k}^{\text{TT}} h_{ik,j}^{\text{TT}} \bar{\phi}_{(2)} - \frac{1}{4} \frac{d-10}{d-2} (h_{ij,k}^{\text{TT}})^2 \bar{\phi}_{(2)} \right) \\ & + (\text{td}). \end{aligned} \quad (43e)$$

By virtue of (40) the post-Newtonian Hamiltonians follow from an integration of the right hand sides of these equations. The last equation leads to the formal 3PN ADM Hamiltonian.

3.3 Momentum Constraint

Also the momentum constraint has to be expanded in a post-Newtonian manner. First of all one has to write the covariant divergence in a more explicit form. For convenience it is also useful to write as much terms as possible in terms of divergences of a traceless symmetric tensor, see [77]. Notice that we also did not remove the trace parts of the field momentum in

$$\begin{aligned}\tilde{\pi}_{,j}^{ij} = & -8\pi G^{(d)} \mathcal{H}_i^{\text{matter}} + \left[\left(1 - (1 + \bar{\phi})^{4/(d-2)} \right) (\tilde{\pi}^{ij} + \pi_{\text{TT}}^{ij}) + V^k (h_{kj,i}^{\text{TT}} + h_{ik,j}^{\text{TT}} - h_{ij,k}^{\text{TT}}) \right. \\ & \left. - \left(1 - \frac{2}{d} \right) V_{,k}^k h_{ij}^{\text{TT}} \right]_{,j} - \Delta(h_{ik}^{\text{TT}} V^k) + \frac{1}{2} h_{\ell j,i}^{\text{TT}} \pi_{\text{TT}}^{\ell j} - (h_{ik}^{\text{TT}} \pi_{\text{TT}}^{kj})_{,j} \\ & + \frac{1}{2} h_{\ell j,i}^{\text{TT}} \tilde{\pi}^{\ell j} - (h_{ik}^{\text{TT}} \tilde{\pi}^{kj})_{,j} - \frac{4}{d-2} (1 + \bar{\phi})^{(6-d)/(d-2)} \left(\tilde{\pi}^{\ell i} \bar{\phi}_{,\ell} - \frac{1}{2} \tilde{\pi}^{\ell\ell} \bar{\phi}_{,i} \right),\end{aligned}\quad (44)$$

as the tracelessness condition may be violated for the non-canonical field momentum due to the spin. The post-Newtonian expansion is necessary for the later integrations by parts and to obtain the necessary field solutions for the momentum type fields. The expansion reads

$$\tilde{\pi}_{(3),j}^{ij} = -8\pi G^{(d)} \mathcal{H}_{i(3)}^{\text{matter}}, \quad (45a)$$

$$\tilde{\pi}_{(5),j}^{ij} = -8\pi G^{(d)} \mathcal{H}_{i(5)}^{\text{matter}} + \left[-\frac{4}{d-2} \tilde{\pi}_{(3)}^{ij} \bar{\phi}_{(2)} \right]_{,j}, \quad (45b)$$

$$\begin{aligned}\tilde{\pi}_{(7),j}^{ij} = & -8\pi G^{(d)} \mathcal{H}_{i(7)}^{\text{matter}} + \left[-\frac{4}{d-2} (\pi_{\text{TT}}^{ij} + \tilde{\pi}_{(5)}^{ij}) \bar{\phi}_{(2)} + \frac{2(d-6)}{(d-2)^2} \tilde{\pi}_{(3)}^{ij} \bar{\phi}_{(2)}^2 - \frac{4}{d-2} \tilde{\pi}_{(3)}^{ij} \bar{\phi}_{(4)} \right. \\ & \left. - \left(1 - \frac{2}{d} \right) h_{ij}^{\text{TT}} V_{(3),k}^k + V_{(3)}^k \left(h_{jk,i}^{\text{TT}} + h_{ik,j}^{\text{TT}} - h_{ij,k}^{\text{TT}} \right) \right]_{,j} - \Delta(h_{ik}^{\text{TT}} V_{(3)}^k).\end{aligned}\quad (45c)$$

Notice that we did not insert the expansion for h_{ij}^{TT} and π_{TT}^{ij} since this is only possible after their evolution were obtained and solved order by order later on.

3.4 Integration by Parts

Due to the complicated structure of $-\Delta\phi_{(10)}/(16\pi G^{(d)})$ in particular the appearance of $\bar{\phi}_{(8)}$, $\bar{\phi}_{(6)}$ and $\tilde{\pi}_{(5)}^{ij}$, $\tilde{\pi}_{(7)}^{ij}$ it is necessary to simplify the integral over the right hand side of (43e). Some of the mentioned fields are not even known in $d = 3$ dimensions. The best way to remove them is to integrate by parts certain terms and afterwards use lower order Hamilton and momentum constraints.

In the used dimensional regularization one may always neglect boundary terms if the integrands are not UV- and IR-divergent simultaneously. These more subtle terms occur at 4PN point-mass calculations for the first time. In the present calculation we always neglected boundary terms in the integrations by parts.

The parts of the Hamilton constraint coming from the expanded Ricci scalar in the conformal approximation have always a structure where a power of the post-Newtonian potential is coupled to a matter source of the Hamilton constraint. These terms can be simplified by inserting the lower order Hamilton constraint for the source and shift the emerging Laplacian to the coupled post-Newtonian potential via integrating by parts twice. This procedure can be used to eliminate $\bar{\phi}_{(8)}$ and $\bar{\phi}_{(6)}$ and the appropriate calculations are given by

$$\begin{aligned}-\bar{\phi}_{(8)} \mathcal{H}_{(2)}^{\text{matter}} = & -\bar{\phi}_{(2)} \mathcal{H}_{(8)}^{\text{matter}} + \bar{\phi}_{(2)}^2 \mathcal{H}_{(6)}^{\text{matter}} + (\bar{\phi}_{(4)} \bar{\phi}_{(2)} - \bar{\phi}_{(2)}^3) \mathcal{H}_{(4)}^{\text{matter}} + (\bar{\phi}_{(6)} \bar{\phi}_{(2)} \\ & - 2\bar{\phi}_{(2)}^2 \bar{\phi}_{(4)} + \bar{\phi}_{(2)}^4) \mathcal{H}_{(2)}^{\text{matter}} - \frac{1}{16\pi G^{(d)}} \left\{ -\frac{3d-10}{d-2} \bar{\phi}_{(2)}^2 (\tilde{\pi}_{(3)}^{ij})^2 \right. \\ & + 2\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{ij} (\tilde{\pi}_{(5)}^{ij} + \pi_{\text{TT}}^{ij}) - \frac{4}{d-2} h_{ij}^{\text{TT}} \bar{\phi}_{(2),i} \bar{\phi}_{(2),j} \bar{\phi}_{(2)} + \frac{1}{4} \bar{\phi}_{(2)} (h_{ij,k}^{\text{TT}})^2 \\ & \left. + \frac{16}{d-2} (\bar{\phi}_{(2)} \bar{\phi}_{(2),i} h_{ij}^{\text{TT}})_{,j} \bar{\phi}_{(2)} - 4(\bar{\phi}_{(4)} h_{ij}^{\text{TT}})_{,ij} \bar{\phi}_{(2)} - \frac{1}{2} \bar{\phi}_{(2)} \Delta(h_{ij}^{\text{TT}})^2 \right\}\end{aligned}$$

$$+ \frac{1}{2} \bar{\phi}_{(2)} (h_{ik}^{\text{TT}} h_{jk}^{\text{TT}})_{,ij} \Big\} + (\text{td}) , \quad (46)$$

$$\begin{aligned} -\bar{\phi}_{(6)} \mathcal{H}_{(4)}^{\text{matter}} &= -\bar{\phi}_{(42)} \mathcal{H}_{(6)}^{\text{matter}} + \bar{\phi}_{(2)} \bar{\phi}_{(42)} \mathcal{H}_{(4)}^{\text{matter}} + (\bar{\phi}_{(4)} \bar{\phi}_{(42)} - \bar{\phi}_{(2)}^2 \bar{\phi}_{(42)}) \mathcal{H}_{(2)}^{\text{matter}} \\ &\quad - \frac{1}{16\pi G^{(d)}} \left\{ \bar{\phi}_{(42)} (\tilde{\pi}_{(3)}^{ij})^2 - 4\bar{\phi}_{(42)} (\bar{\phi}_{(2)} h_{ij}^{\text{TT}})_{,ij} \right\} + (\text{td}) , \end{aligned} \quad (47)$$

$$\begin{aligned} 3\bar{\phi}_{(6)} \bar{\phi}_{(2)} \mathcal{H}_{(2)}^{\text{matter}} &= \left\{ -3\mathcal{H}_{(6)}^{\text{matter}} + 3\bar{\phi}_{(2)} \mathcal{H}_{(4)}^{\text{matter}} - (-3\bar{\phi}_{(4)} + 3\bar{\phi}_{(2)}^2) \mathcal{H}_{(2)}^{\text{matter}} \right. \\ &\quad \left. - \frac{1}{16\pi G^{(d)}} \left[3(\tilde{\pi}_{(3)}^{ij})^2 - 12\bar{\phi}_{(2),ij} h_{ij}^{\text{TT}} \right] \right\} \bar{\phi}_{(40)} + (\text{td}) . \end{aligned} \quad (48)$$

The fields $\bar{\phi}_{(42)}$ and $\bar{\phi}_{(40)}$ are the momentum dependent and the momentum independent part of $\bar{\phi}_{(4)}$ given by $\bar{\phi}_{(42)} = -16\pi G^{(d)} \frac{d-2}{4(d-1)} \Delta^{-1} \mathcal{H}_{(4)}^{\text{matter}}$ and $\bar{\phi}_{(40)} = -16\pi G^{(d)} \frac{d-2}{4(d-1)} \Delta^{-1} (-\bar{\phi}_{(2)} \mathcal{H}_{(2)}^{\text{matter}})$. It is possible to simplify (46) once more using

$$-\frac{1}{16\pi G^{(d)}} \left(-\frac{1}{2} \bar{\phi}_{(2)} \Delta (h_{ij}^{\text{TT}})^2 \right) = -\frac{d-2}{8(d-1)} (h_{ij}^{\text{TT}})^2 \mathcal{H}_{(2)}^{\text{matter}} + (\text{td}) . \quad (49)$$

In the Hamilton constraint there are also several terms coupling different orders of the longitudinal field momentum. These terms can also be simplified by inserting the decomposition (34), removing the derivatives from the vector potential V^i via an integration by parts and inserting the lower order momentum constraints (45) into the divergencies of the longitudinal field momentum. This procedure is used in order to eliminate $\tilde{\pi}_{(5)}^{ij}$ and $\tilde{\pi}_{(7)}^{ij}$. The respective integrations by parts are given by

$$\begin{aligned} -\frac{1}{16\pi G^{(d)}} \left(-2\tilde{\pi}_{(7)}^{ij} \tilde{\pi}_{(3)}^{ij} \right) &= 2V_{(3)}^i \mathcal{H}_{i(7)}^{\text{matter}} - \frac{1}{16\pi G^{(d)}} \left\{ -2h_{ij}^{\text{TT}} \left[-2V_{(3),k}^j \tilde{\pi}_{(3)}^{ik} + (V_{(3)}^k \tilde{\pi}_{(3)}^{ij})_{,k} \right] \right. \\ &\quad + \frac{8}{d-2} \tilde{\pi}_{(3)}^{ij} (\pi_{\text{TT}}^{ij} + \tilde{\pi}_{(5)}^{ij}) \bar{\phi}_{(2)} - 4 \frac{d-6}{(d-2)^2} (\tilde{\pi}_{(3)}^{ij})^2 \bar{\phi}_{(2)}^2 \\ &\quad \left. + \frac{8}{d-2} (\tilde{\pi}_{(3)}^{ij})^2 \bar{\phi}_{(4)} \right\} + (\text{td}) . \end{aligned} \quad (50)$$

Last but not least the $(\tilde{\pi}_{(5)}^{ij})^2$ integration by parts is given by

$$-\frac{1}{16\pi G^{(d)}} \left(-(\tilde{\pi}_{(5)}^{ij})^2 \right) = V_{(5)}^i \mathcal{H}_{i(5)}^{\text{matter}} - \frac{1}{16\pi G^{(d)}} \left\{ \frac{4}{d-2} \tilde{\pi}_{(3)}^{ij} \tilde{\pi}_{(5)}^{ij} \bar{\phi}_{(2)} \right\} + (\text{td}) . \quad (51)$$

Although it is good to express field integrals in terms of sources and fields, the $V_{(5)}^i$ potential is still very complicated. Thus we try to express $(\tilde{\pi}_{(5)}^{ij})^2$ in yet another way. From the transverse-traceless projection of an arbitrary second rank tensor field A_{ij} , namely

$$\delta_{kl}^{\text{TT}ij} A_{kl} = \left(\delta_{i(k} \delta_{\ell)j} - \delta_{kl}^{\text{LT}ij} - \delta_{kl}^{\text{Tr}ij} \right) A_{kl} , \quad (52)$$

$$\begin{aligned} &= A_{ij} - \Delta^{-1} \left((A_{j\ell})_{,\ell i} + (A_{i\ell})_{,\ell j} - \frac{1}{d-1} \mathcal{D}_{d-1}^{ij} (A_{kl})_{,kl} \right) \\ &\quad - \frac{1}{d-1} (\delta_{ij} - \partial_i \partial_j \Delta^{-1}) \delta_{kl} A_{kl} , \end{aligned} \quad (53)$$

one can get the transverse-traceless projection $(\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{kl})$ is traceless)

$$\delta_{kl}^{\text{TT}ij} (\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{kl}) = \bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{ij} + \frac{d-2}{4} \left\{ \tilde{\pi}_{(5)}^{ij} + \tilde{\pi}_{(5)1}^{ij} \right\} , \quad (54)$$

via the momentum constraint (45). Here

$$\tilde{\pi}_{(5)1}^{ij} = \tilde{\pi}_{(5)1,j}^i + \tilde{\pi}_{(5)1,i}^j - \frac{1}{d-1} \mathcal{D}_{d-1}^{ij} \tilde{\pi}_{(5)1,k}^k , \quad (55)$$

where

$$\tilde{\pi}_{(5)1}^i = 8\pi G^{(d)} \Delta^{-1} \mathcal{H}_{i(5)}^{\text{matter}}. \quad (56)$$

Notice that $\tilde{\pi}_{(5)1}^i$ is defined with a different sign than the usual vector potentials used throughout this article. Now we can express $\tilde{\pi}_{(5)}^{ij}$ in terms of $\tilde{\pi}_{(5)1}^i$, $\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{ij}$ and the transverse-traceless projection (54). From these considerations it follows that $(\tilde{\pi}_{(5)}^{ij})^2$ is given by

$$\begin{aligned} -\frac{1}{16\pi G^{(d)}} \left(-(\tilde{\pi}_{(5)}^{ij})^2 \right) &= -\frac{1}{16\pi G^{(d)}} \left\{ \frac{16}{(d-2)^2} \left[\left(\delta_{k\ell}^{\text{TT}ij} (\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{k\ell}) \right)^2 - \bar{\phi}_{(2)}^2 (\tilde{\pi}_{(3)}^{ij})^2 \right] \right. \\ &\quad \left. - \frac{8}{d-2} \bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{ij} \tilde{\pi}_{(5)1}^i - (\tilde{\pi}_{(5)1}^i)^2 \right\} + (\text{td}). \end{aligned} \quad (57)$$

Some of the results above were partially checked by comparison with [77].

3.5 The formal 3PN ADM Hamiltonian

Performing the mentioned integration by parts leads to the following formal 3PN ADM Hamiltonian which can also be compared to [77],

$$\begin{aligned} H_{3\text{PN}} &= \int d^d x \left[\mathcal{H}_{(10)}^{\text{matter}} - 2\bar{\phi}_{(2)} \mathcal{H}_{(8)}^{\text{matter}} + (-\bar{S}_{(4)} - 4\bar{\phi}_{(4)} + 2\bar{\phi}_{(2)}^2) \mathcal{H}_{(6)}^{\text{matter}} + (6\bar{\phi}_{(2)} \bar{\phi}_{(4)} + \bar{S}_{(4)} \bar{\phi}_{(2)}) \right. \\ &\quad \left. - 2\bar{\phi}_{(2)}^3 \mathcal{H}_{(4)}^{\text{matter}} + \left(4\bar{\phi}_{(4)}^2 + \bar{S}_{(4)} \bar{\phi}_{(4)} - 8\bar{\phi}_{(4)} \bar{\phi}_{(2)}^2 - \bar{S}_{(4)} \bar{\phi}_{(2)}^2 + 2\bar{\phi}_{(2)}^4 \right. \right. \\ &\quad \left. \left. - \frac{1}{4(d-1)} (h_{ij}^{\text{TT}})^2 \right) \mathcal{H}_{(2)}^{\text{matter}} + 2V_{(3)}^i \mathcal{H}_{i(7)}^{\text{matter}} - \frac{1}{16\pi G^{(d)}} \left(-(\tilde{\pi}_{(5)1}^i)^2 + 2\frac{3d-4}{d-2} \bar{\phi}_{(4)} (\tilde{\pi}_{(3)}^{ij})^2 \right. \right. \\ &\quad \left. \left. + \bar{S}_{(4)} (\tilde{\pi}_{(3)}^{ij})^2 - \frac{(3d-4)(3d-2)}{(d-2)^2} (\tilde{\pi}_{(3)}^{ij})^2 \bar{\phi}_{(2)}^2 - (\pi_{\text{TT}}^{ij})^2 + 8\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{ij} \pi_{\text{TT}}^{ij} \right. \right. \\ &\quad \left. \left. - 2h_{ij}^{\text{TT}} \left[-2V_{(3),k}^j \tilde{\pi}_{(3)}^{ik} + (V_{(3)}^k \tilde{\pi}_{(3)}^{ij})_{,k} + \tilde{\pi}_{(3)}^{ik} \tilde{\pi}_{(3)}^{jk} - 4\frac{2d-3}{d-2} \bar{\phi}_{(2),i} \bar{\phi}_{(4),j} \right. \right. \right. \\ &\quad \left. \left. - 4\frac{3d-4}{(d-2)^2} \bar{\phi}_{(2)} \bar{\phi}_{(2),i} \bar{\phi}_{(2),j} - 2\bar{S}_{(4),i} \bar{\phi}_{(2),j} \right] - 8\frac{d-1}{d-2} \bar{\phi}_{(2)} \tilde{\pi}_{(5)1}^i \tilde{\pi}_{(3)}^{ij} \right. \right. \\ &\quad \left. \left. + 16\frac{2d-3}{(d-2)^2} \left(\delta_{k\ell}^{\text{TT}ij} (\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{k\ell}) \right)^2 + \frac{2}{d-2} (h_{ij,k}^{\text{TT}})^2 \bar{\phi}_{(2)} \right) \right]. \end{aligned} \quad (58)$$

We changed all occurrences of $\bar{\phi}_{(4)2}$ to $\bar{S}_{(4)} = -2\bar{\phi}_{(4)2}$ to gain a result which is comparable to [31]. Note that, since we did not expand h_{ij}^{TT} and $\hat{\pi}_{\text{TT}}^{ij}$, we also need some formal 2PN terms which may contribute to the 3PN kinetic energy or the 3PN interaction Hamiltonian after expanding h_{ij}^{TT} ,

$$H_{2\text{PN}} = \int d^d x \left[\mathcal{H}_{(8)\text{TT}}^{\text{matter}} + \dots - \frac{1}{16\pi G^{(d)}} \left(\dots + \frac{4(d-1)}{d-2} \bar{\phi}_{(2),i} \bar{\phi}_{(2),j} h_{ij}^{\text{TT}} - \frac{1}{4} (h_{ij,k}^{\text{TT}})^2 \right) \right]. \quad (59)$$

Now we can split up the Hamiltonian into a kinetic part for the reduced canonical field variables $[h_{ij}^{\text{TT}}$ and $\hat{\pi}_{\text{TT}}^{ij}$; after inserting (37)], an interaction part between these canonical fields and constraint fields, and a part independent of the canonical fields, i.e.,

$$H_{\text{ADM}} = H_{\text{ADM}}^{\text{int}} + H_{\text{ADM}}^{\text{non-TT}} + \frac{1}{16\pi G^{(d)}} \int d^d x \left[\frac{1}{4} (h_{ij,k}^{\text{TT}})^2 + (\hat{\pi}_{\text{TT}}^{ij})^2 \right], \quad (60)$$

with

$$H_{\text{ADM}}^{\text{int}} = \frac{1}{16\pi G^{(d)}} \int d^d x \left\{ (B_{(4)ij} + \hat{B}_{(6)ij}) h_{ij}^{\text{TT}} - \frac{4\pi G^{(d)}}{d-1} (h_{ij}^{\text{TT}})^2 \mathcal{H}_{(2)}^{\text{matter}} + \hat{\pi}_{\text{TT}}^{ij} C_{ij} \right\}$$

$$- \frac{2}{d-2} \bar{\phi}_{(2)} (h_{ij,k}^{\text{TT}})^2 \Big\}, \quad (61)$$

$$H_{\text{ADM}}^{\text{non-TT}} = H_{\text{ADM}} - H_{\text{ADM}}^{\text{int}} - \frac{1}{16\pi G^{(d)}} \int d^d x \left[\frac{1}{4} (h_{ij,k}^{\text{TT}})^2 + (\hat{\pi}_{\text{TT}}^{ij})^2 \right], \quad (62)$$

and

$$B_{(4)ij} = 16\pi G^{(d)} \frac{\delta}{\delta h_{ij}^{\text{TT}}} \int d^d x \mathcal{H}_{(8)}^{\text{matter}} - \frac{4(d-1)}{d-2} \bar{\phi}_{(2),i} \bar{\phi}_{(2),j}, \quad (63)$$

$$\begin{aligned} \hat{B}_{(6)ij} = 16\pi G^{(d)} \frac{\delta}{\delta h_{ij}^{\text{TT}}} \int d^d x & \left[\mathcal{H}_{(10)}^{\text{matter}} - 2\bar{\phi}_{(2)} \mathcal{H}_{(8)}^{\text{matter}} + 2V_{(3)}^k \mathcal{H}_{k(7)}^{\text{matter}} \right] + 2 \left[-2V_{(3),k}^j \tilde{\pi}_{(3)}^{ik} \right. \\ & + (V_{(3)}^k \tilde{\pi}_{(3)}^{ij})_{,k} + \tilde{\pi}_{(3)}^{ik} \tilde{\pi}_{(3)}^{jk} - 4 \frac{2d-3}{d-2} \bar{\phi}_{(2),i} \bar{\phi}_{(4),j} + 4 \frac{3d-4}{(d-2)^2} \bar{\phi}_{(2)} \bar{\phi}_{(2),i} \bar{\phi}_{(2),j} \\ & \left. - 2\bar{S}_{(4),i} \bar{\phi}_{(2),j} \right], \end{aligned} \quad (64)$$

$$C_{ij} = -2\pi_{\text{matter}}^{ij} - 8\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{ij}. \quad (65)$$

H_{ADM} consists of all Hamiltonians starting from the rest mass contribution up to $H_{3\text{PN}}$. The $B_{(4)ij}$ part in $H_{\text{ADM}}^{\text{int}}$ comes from the formal 2PN Hamiltonian shown in (59). $H_{\text{ADM}}^{\text{non-TT}}$ can be obtained by removing all h_{ij}^{TT} and $\hat{\pi}_{\text{TT}}^{ij}$ parts in (59) and (58) respectively. Notice that the relevant source terms in the expressions for $B_{(4)ij}$ and $\hat{B}_{(6)ij}$ are at most linear in h_{ij}^{TT} . This allowed us to single out these contributions by a functional derivative.

Now we arrived at a point mentioned in Sect. 2.4, namely where we are able to eliminate the constraint fields using lower order Hamilton and momentum constraints, but where the dynamical field degrees of freedom are still in the Hamiltonian. The elimination of these and further simplifications of the calculation process are subject of the following section.

4 Routhian and Application of Wave Equation

To obtain a fully reduced matter only Hamiltonian, we have to remove the dynamical degrees of freedom h_{ij}^{TT} and $\hat{\pi}_{\text{TT}}^{ij}$ by solving the appropriate equations of motion and inserting their solutions into H_{ADM} . However there are some subtleties in this procedure which will be discussed in Sect. 4.2.

From Hamilton's equations

$$\frac{1}{16\pi G^{(d)}} \frac{\partial h_{ij}^{\text{TT}}}{\partial t} = \delta_{kl}^{\text{TT}ij} \frac{\delta H_{\text{ADM}}}{\delta \hat{\pi}_{\text{TT}}^{kl}}, \quad (66)$$

$$\frac{1}{16\pi G^{(d)}} \frac{\partial \hat{\pi}_{\text{TT}}^{ij}}{\partial t} = -\delta_{kl}^{\text{TT}ij} \frac{\delta H_{\text{ADM}}}{\delta h_{kl}^{\text{TT}}}, \quad (67)$$

and the split of the ADM Hamiltonian (60) one gets the appropriate wave equations

$$\square h_{ij}^{\text{TT}} = 16\pi G^{(d)} \delta_{kl}^{\text{TT}ij} \left[2 \frac{\delta H_{\text{ADM}}^{\text{int}}}{\delta h_{kl}^{\text{TT}}} - \frac{\partial}{\partial t} \frac{\delta H_{\text{ADM}}^{\text{int}}}{\delta \hat{\pi}_{\text{TT}}^{kl}} \right], \quad (68)$$

$$\hat{\pi}_{\text{TT}}^{ij} = \frac{1}{2} \left[\dot{h}_{ij}^{\text{TT}} - 16\pi G^{(d)} \delta_{kl}^{\text{TT}ij} \frac{\delta H_{\text{ADM}}^{\text{int}}}{\delta \hat{\pi}_{\text{TT}}^{kl}} \right], \quad (69)$$

for the dynamical degrees of freedom of the gravitational field (here $\square = \Delta - c^{-2} \partial_t^2$; remember that we use a $(d+1)$ -metric with signature $d-1$). In order to receive the full wave equations one has to perform the variational derivative of the interaction Hamiltonian,

$$\square h_{ij}^{\text{TT}} = \delta_{kl}^{\text{TT}ij} \left\{ 2(B_{(4)kl} + \hat{B}_{(6)kl}) - \frac{16\pi G^{(d)}}{d-1} \mathcal{H}_{(2)}^{\text{matter}} h_{kl}^{\text{TT}} + \frac{8}{d-2} (\bar{\phi}_{(2)} h_{kl,m}^{\text{TT}})_{,m} - \frac{\partial}{\partial t} C_{kl} \right\}, \quad (70)$$

$$\hat{\pi}_{\text{TT}}^{ij} = \frac{1}{2} h_{ij}^{\text{TT}} - \frac{1}{2} \delta_{kl}^{\text{TT}ij} C_{kl}. \quad (71)$$

4.1 Near-Zone Expansion

The wave equation

$$\square h = f, \quad (72)$$

[which represents the components of (70)] for a source f has several solutions depending on the boundary conditions. In field theory mostly the retarded one is used. At the order considered in the present article the time symmetric (i.e. conservative) solution is sufficient.³

The appropriate solution derives from a near-zone expansion, which is a formal expansion in c^{-1} . (see e.g. [181–184].) More precisely, the near-zone expansion is a series expansion in the small quantity $r/(ct) \ll 1$, which means that the distance from the field point to the appropriate source point is small compared to the gravitational wavelength. So the near-zone expansion may be used if retardation effects are negligible. Consider the Feynman propagator for a massless particle (which corresponds to the Green's function of the wave equation)

$$G_F(\mathbf{x}, t) = -\frac{1}{2\pi} \lim_{\varepsilon \rightarrow 0} \int dk_0 \frac{1}{(2\pi)^d} \int d^d k \frac{e^{-i(k_0 x^0 - (\mathbf{k} \cdot \mathbf{x}))}}{k^2 - k_0^2 - i\varepsilon}. \quad (73)$$

For the reader's convenience we reintroduce the powers of c in the following expressions. This means $k_0 = \omega/c$ and $x^0 = ct$. As it was argued above and in e.g. [185, 186] the condition $r/(ct) \ll 1$ corresponds to $k_0/k \ll 1$ which gives rise to the so-called potential gravitons in contrast to the radiation gravitons for which $k_0^2 \approx k^2$. For the radiation gravitons the $i\varepsilon$ term becomes important, but for the potential ones it could be neglected and the Fourier amplitude in the propagator is expandable in k_0/k , namely

$$G_F(\mathbf{x}, t) \stackrel{\text{NZ}}{=} -\frac{1}{2\pi c} \int d\omega e^{-i\omega t} \sum_{n=0}^{\infty} \frac{\omega^{2n}}{c^{2n}} \frac{1}{(2\pi)^d} \int d^d k \frac{e^{i(\mathbf{k} \cdot \mathbf{x})}}{k^{2(n+1)}}. \quad (74)$$

This means that a near-zone expansion of the (time symmetric) Feynman propagator cannot contain any contributions leading to radiative losses. The d -dimensional k -space integral in (74) has to be performed by using

$$\int d^d k \frac{e^{i(\mathbf{k} \cdot \mathbf{x})}}{k^{2\alpha}} = \frac{1}{(4\pi)^{d/2}} \frac{\Gamma(\frac{d}{2} - \alpha)}{\Gamma(\alpha)} \left(\frac{r^2}{4}\right)^{\alpha-d/2}, \quad (75)$$

which can be obtained by dimensional regularization (the source of the wave lies at the origin of the coordinate system). This result can be reformulated in inverse Laplacians on a delta-type source by iterating (167) and (168) in the Appendix,

$$\frac{1}{(2\pi)^d} \int d^d k \frac{e^{i(\mathbf{k} \cdot \mathbf{x})}}{k^{2(n+1)}} = -\frac{\Gamma(2+n-\frac{d}{2}) \Gamma(\frac{d}{2}-1-n)}{\Gamma(2-\frac{d}{2}) \Gamma(\frac{d}{2}-1)} (\Delta^{-1})^{n+1} \delta. \quad (76)$$

For $d \notin 2\mathbb{Z}$ (i.e. no odd dimensional spacetime) one can simplify the Gamma functions further by using the identity $\Gamma(1-z)\Gamma(z) = \pi/\sin(\pi z)$, $z \notin \mathbb{Z}$ which leads to

$$\frac{1}{(2\pi)^d} \int d^d k \frac{e^{i(\mathbf{k} \cdot \mathbf{x})}}{k^{2(n+1)}} = -(-1)^n (\Delta^{-1})^{n+1} \delta, \quad (77)$$

³At the considered order $h_{(5)ij}^{\text{TT}}$ can be neglected since the linear terms are not time symmetric and the quadratic terms arise only due to $(h_{(5)ij,k}^{\text{TT}})^2$ which is zero in $d = 3$ because $h_{(5)ij}^{\text{TT}}$ is only a function of t (there are no explicit terms in the source at c^{-5} -order); $h_{(7)ij}^{\text{TT}}$ is of too high order to appear at formal 3PN order, see [32, 59, 77]. We therefore neglected $h_{(5)ij}^{\text{TT}}$ and $h_{(7)ij}^{\text{TT}}$ in (41e) and hence there are no contributions to (58) or (59).

and therefore after performing the ω Fourier transform to

$$G_F(\mathbf{x}, t) \stackrel{\text{NZ}}{=} \sum_{n=0}^{\infty} \left(\frac{1}{c} \frac{\partial}{\partial t} \right)^{2n} (\Delta^{-1})^{n+1} \delta_{(d+1)}, \quad (78)$$

for the Feynman propagator. This result could immediately be used to write down the near-zone expanded solution of the wave equation

$$h = \square^{-1} f \stackrel{\text{NZ}}{=} \Delta^{-1} \sum_{n=0}^{\infty} (\Delta^{-1})^n \left(\frac{1}{c} \frac{\partial}{\partial t} \right)^{2n} f. \quad (79)$$

Notice that a near-zone expanded field in general does not converge at spatial infinity.

Now we are able to derive the solutions for the transverse-traceless part of the metric at a certain post-Newtonian order in the near-zone.

4.2 Routhian

Before we can insert the wave equation (70) and its solution [see (79), and for a more explicit form see Sect. 6.2], we have to transform the ADM Hamiltonian into a Routhian, i.e. a Lagrangian in h_{ij}^{TT} and $\hat{\pi}_{\text{TT}}^{ij}$ and a Hamiltonian in the particle degrees of freedom,

$$R[h_{ij}^{\text{TT}}, \dot{h}_{ij}^{\text{TT}}] = H_{\text{ADM}} - \frac{1}{16\pi G^{(d)}} \int d^d x \hat{\pi}_{\text{TT}}^{ij} \dot{h}_{ij}^{\text{TT}}. \quad (80)$$

This is necessary because otherwise the equation of motion, e.g., for $\hat{\mathbf{z}}_a$, following from the Hamiltonian (for simplicity we omit the spin variables here)

$$H_{\text{ADM}}(\hat{\mathbf{z}}_a, \hat{\mathbf{p}}_a, h_{ij}^{\text{TT}}(\hat{\mathbf{z}}_b, \hat{\mathbf{p}}_b), \hat{\pi}_{\text{TT}}^{ij}(\hat{\mathbf{z}}_b, \hat{\mathbf{p}}_b)), \quad (81)$$

using the Poisson brackets would be

$$\dot{\hat{\mathbf{z}}}_a^k = \frac{\partial H_{\text{ADM}}}{\partial \hat{p}_{a k}} + \frac{\delta H_{\text{ADM}}}{\delta h_{ij}^{\text{TT}}} \frac{\partial h_{ij}^{\text{TT}}}{\partial \hat{p}_{a k}} + \frac{\delta H_{\text{ADM}}}{\delta \hat{\pi}_{\text{TT}}^{ij}} \frac{\partial \hat{\pi}_{\text{TT}}^{ij}}{\partial \hat{p}_{a k}}. \quad (82)$$

This equation is obviously wrong, because h_{ij}^{TT} and $\hat{\pi}_{\text{TT}}^{ij}$ are dynamical degrees of freedom and may not lead to additional terms in the equations of motion, also when their solutions are inserted. On the other hand using $R(\hat{\mathbf{z}}_a, \hat{\mathbf{p}}_a, h_{ij}^{\text{TT}}(\hat{\mathbf{z}}_b, \hat{\mathbf{p}}_b), \dot{h}_{ij}^{\text{TT}}(\hat{\mathbf{z}}_b, \dot{\hat{\mathbf{z}}}_b, \hat{\mathbf{p}}_b, \dot{\hat{\mathbf{p}}}_b))$ one has for the same equation of motion

$$\dot{\hat{\mathbf{z}}}_a^k = \frac{\partial R}{\partial \hat{p}_{a k}} + \underbrace{\frac{\delta R}{\delta h_{ij}^{\text{TT}}} \frac{\partial h_{ij}^{\text{TT}}}{\partial \hat{p}_{a k}}}_{=0}, \quad (83)$$

which has no additional terms coming from the chain rule, because they vanish due to h_{ij}^{TT} fulfilling the equations of motion in the appropriate approximation. Hence one can obtain the equation of motion in the usual way and one does not have to keep track of the field insertions. This is analogous to the construction of a Fokker action, see, e.g., [37] and references therein.

A Fokker-like construction of a matter-only Lagrangian (or Routhian) can not account for dissipative effects, see [187] for a discussion. However, dissipative Hamiltonians in the ADM formalism can be constructed in the following way [32, 59]: The matter variables entering the solution of h_{ij}^{TT} and $\hat{\pi}_{\text{TT}}^{ij}$ are substituted by new (“primed”) variables and thus the binary Hamiltonian now contains four types of canonical matter variables. This procedure prevents occurrence of wrong contributions in the equations of motion, too. The primed variables will be treated as explicitly time dependent and lead to an explicitly time dependent Hamiltonian. After calculating the equations of motion for canonical positions $\hat{\mathbf{z}}_a$ and momenta $\hat{\mathbf{p}}_a$, the primed variables are again identified with the old ones which makes another regularization procedure necessary [32]. Another approach to construct an action principle for dissipative systems was suggested in [187] recently.

4.3 Reduction of the Routhian using the Wave Equation

The part of the Routhian labeled as TT contains $H_{\text{ADM}}^{\text{int}}$, the field kinetic part of the Hamiltonian and the part coming from the Legendre transform (and thus only terms coming from the transverse-traceless degrees of freedom),

$$R^{\text{TT}} = \frac{1}{16\pi G^{(d)}} \int d^d x \left[\left(B_{(4)ij} + \hat{B}_{(6)ij} \right) h_{ij}^{\text{TT}} - \frac{4\pi G^{(d)}}{d-1} (h_{ij}^{\text{TT}})^2 \mathcal{H}_{(2)}^{\text{matter}} + \hat{\pi}_{\text{TT}}^{ij} C_{ij} - \frac{2}{d-2} \bar{\phi}_{(2)} (h_{ij,k}^{\text{TT}})^2 + \frac{1}{4} (h_{ij,k}^{\text{TT}})^2 + (\hat{\pi}_{\text{TT}}^{ij})^2 - \dot{h}_{ij}^{\text{TT}} \hat{\pi}_{\text{TT}}^{ij} \right]. \quad (84)$$

Inserting $\square h_{ij}^{\text{TT}}$ and $\hat{\pi}_{\text{TT}}^{ij}$ from (70) and (71), or

$$h_{ij}^{\text{TT}} (B_{(4)ij} + \hat{B}_{(6)ij}) = \frac{1}{2} h_{ij}^{\text{TT}} \square h_{ij}^{\text{TT}} + \frac{8\pi G^{(d)}}{d-1} (h_{ij}^{\text{TT}})^2 \mathcal{H}_{(2)}^{\text{matter}} - \frac{4}{d-2} h_{ij}^{\text{TT}} (\bar{\phi}_{(2)} h_{ij,k}^{\text{TT}})_{,k} + \frac{1}{2} h_{ij}^{\text{TT}} \frac{\partial}{\partial t} C_{ij} + (\text{td}), \quad (85)$$

leads to

$$R^{\text{TT}} = \frac{1}{16\pi G^{(d)}} \int d^d x \left[\frac{1}{4} h_{ij}^{\text{TT}} \square h_{ij}^{\text{TT}} + \frac{4\pi G^{(d)}}{d-1} (h_{ij}^{\text{TT}})^2 \mathcal{H}_{(2)}^{\text{matter}} + \frac{2}{d-2} \bar{\phi}_{(2)} (h_{ij,k}^{\text{TT}})^2 + \frac{1}{2} \frac{\partial}{\partial t} \left[h_{ij}^{\text{TT}} C_{ij} \right] - \frac{1}{4} C_{ij} \delta_{k\ell}^{\text{TT} ij} C_{k\ell} \right], \quad (86)$$

where the last part will appear in the matter part of the final Routhian, because there is no h_{ij}^{TT} and no $\hat{\pi}_{\text{TT}}^{ij}$ or $\dot{h}_{ij}^{\text{TT}}$ in C_{ij} . Notice that we kept a total time derivative here. If we would drop it the $\dot{C}_{ij}$ terms would not cancel in the next step. These terms are not impossible to handle, but it is advised to remove them to simplify the calculation.

4.4 Insertion of the Near-Zone Wave Equation for Further Simplification

Now we need to split up the first expression in the TT part of the Routhian (86). The near-zone expansion of the transverse-traceless part of the metric $h_{ij}^{\text{TT}} = h_{(4)ij}^{\text{TT}} + h_{(6)ij}^{\text{TT}} + \dots$, which is explained in detail in Sect. 4.1, contributes only via $h_{(4)ij}^{\text{TT}}$ and $h_{(6)ij}^{\text{TT}}$ at 3PN level. This expansion leads to

$$h_{ij}^{\text{TT}} \square h_{ij}^{\text{TT}} = h_{(4)ij}^{\text{TT}} \square h_{(4)ij}^{\text{TT}} + 2h_{(4)ij}^{\text{TT}} \square h_{(6)ij}^{\text{TT}} + (\text{td}) + (\text{ttd}), \quad (87)$$

where (ttd) denotes a total time derivative. Notice that there is a difference between $\square h_{(6)ij}^{\text{TT}}$ and $(\square h_{ij}^{\text{TT}})_{(6)}$ in the near-zone expansion as time derivatives raise the formal order of a field in contrast to spatial derivatives. This is a specific feature of the near-zone expansion, as in the far-zone time and space derivatives are equal in magnitude. These considerations lead to the difference in the following box operations,

$$\square h_{(6)ij}^{\text{TT}} = \Delta h_{(6)ij}^{\text{TT}} - \partial_t^2 h_{(6)ij}^{\text{TT}}, \quad (88)$$

$$(\square h_{ij}^{\text{TT}})_{(6)} = \Delta h_{(6)ij}^{\text{TT}} - \partial_t^2 h_{(4)ij}^{\text{TT}}, \quad (89)$$

where $\partial_t^2 h_{(6)ij}^{\text{TT}}$ is of formal c^{-8} order and hence the total time derivative can be neglected. From this it follows that

$$h_{ij}^{\text{TT}} \square h_{ij}^{\text{TT}} = h_{(4)ij}^{\text{TT}} \Delta h_{(4)ij}^{\text{TT}} - h_{(4)ij}^{\text{TT}} \partial_t^2 h_{(4)ij}^{\text{TT}} + 2h_{(4)ij}^{\text{TT}} \underbrace{\Delta h_{(6)ij}^{\text{TT}}}_{(\square h_{ij}^{\text{TT}})_{(6)} + \partial_t^2 h_{(4)ij}^{\text{TT}}} + (\text{td}), \quad (90)$$

such that

$$(h_{ij}^{\text{TT}} \square h_{ij}^{\text{TT}})_{(10)} = 2h_{(4)ij}^{\text{TT}} (\square h_{ij}^{\text{TT}})_{(6)} + h_{(4)ij}^{\text{TT}} \partial_t^2 h_{(4)ij}^{\text{TT}} + (\text{td}), \quad (91)$$

where $(\square h_{ij}^{\text{TT}})_{(6)}$ is given by the c^{-6} part of Eq. (70). After another integration by parts this immediately leads to (we need only the leading order of C_{ij} which is at c^{-5})

$$R_{3\text{PN}}^{\text{TT}} = \frac{1}{16\pi G^{(d)}} \int d^d x \left[h_{(4)ij}^{\text{TT}} \hat{B}_{(6)ij} - \frac{1}{4} (\dot{h}_{(4)ij}^{\text{TT}})^2 - \frac{4\pi G^{(d)}}{d-1} (h_{(4)ij}^{\text{TT}})^2 \mathcal{H}_{(2)}^{\text{matter}} \right. \\ \left. - \frac{2}{d-2} \bar{\phi}_{(2)} (h_{(4)ij,k}^{\text{TT}})^2 + \frac{1}{2} \dot{h}_{(4)ij}^{\text{TT}} C_{(5)ij} - \frac{1}{4} C_{(5)ij} \delta_{k\ell}^{\text{TT}ij} C_{(5)k\ell} \right]. \quad (92)$$

Rearranging some of the terms into pure matter and pure transverse-traceless parts, one obtains the full final 3PN Routhian,

$$R_{3\text{PN}}^{\text{matter}} = \int d^d x \left[\mathcal{H}_{(10)\text{non-TT}}^{\text{matter}} - 2\bar{\phi}_{(2)} \mathcal{H}_{(8)\text{non-TT}}^{\text{matter}} + (-\bar{S}_{(4)} - 4\bar{\phi}_{(4)} + 2\bar{\phi}_{(2)}^2) \mathcal{H}_{(6)}^{\text{matter}} \right. \\ + (6\bar{\phi}_{(2)}\bar{\phi}_{(4)} + \bar{S}_{(4)}\bar{\phi}_{(2)} - 2\bar{\phi}_{(2)}^3) \mathcal{H}_{(4)}^{\text{matter}} + \left(4\bar{\phi}_{(4)}^2 + \bar{S}_{(4)}\bar{\phi}_{(4)} - 8\bar{\phi}_{(4)}\bar{\phi}_{(2)}^2 - \bar{S}_{(4)}\bar{\phi}_{(2)}^2 \right. \\ \left. + 2\bar{\phi}_{(2)}^4 \right) \mathcal{H}_{(2)}^{\text{matter}} + 2V_{(3)}^i \mathcal{H}_{i(7)\text{non-TT}}^{\text{matter}} - \frac{1}{16\pi G^{(d)}} \left(-(\tilde{\pi}_{(5)1}^{ij})^2 + 2\frac{3d-4}{d-2} \bar{\phi}_{(4)} (\tilde{\pi}_{(3)}^{ij})^2 \right. \\ \left. + \bar{S}_{(4)} (\tilde{\pi}_{(3)}^{ij})^2 - \frac{(3d-4)(3d-2)}{(d-2)^2} (\tilde{\pi}_{(3)}^{ij})^2 \bar{\phi}_{(2)}^2 + \left(\frac{4(d-1)}{d-2} \delta_{k\ell}^{\text{TT}ij} (\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{k\ell}) \right)^2 \right. \\ \left. - 4\frac{2d-3}{d-2} \bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{ij} \tilde{\pi}_{(5)1}^{ij} \right) \right], \quad (93a)$$

$$R_{3\text{PN}}^{\text{TT}} = \frac{1}{16\pi G^{(d)}} \int d^d x \left[h_{(4)ij}^{\text{TT}} \hat{B}_{(6)ij} - \frac{1}{4} (\dot{h}_{(4)ij}^{\text{TT}})^2 - \frac{4\pi G^{(d)}}{d-1} (h_{(4)ij}^{\text{TT}})^2 \mathcal{H}_{(2)}^{\text{matter}} \right. \\ \left. - \frac{2}{d-2} \bar{\phi}_{(2)} (h_{(4)ij,k}^{\text{TT}})^2 - \dot{h}_{(4)ij}^{\text{TT}} \pi_{(5)\text{matter}}^{ij} - 4\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{ij} \dot{h}_{(4)ij}^{\text{TT}} \right], \quad (93b)$$

where $\hat{B}_{(6)ij}$ is given by (64). Note, that one does not need to calculate the $h_{(6)ij}^{\text{TT}}$ field, which is not fully known in closed form. An explicit form of $\hat{B}_{(6)ij}$ with all sources inserted is derived in the next section, cf. (107b).

5 Sources

The construction of the sources linear in spin follows along the lines of [75]. This requires the introduction of a local Lorentz frame⁴ as we want flat space Poisson brackets for the spin. In particular we also apply the Schwinger time gauge for the $(d+1)$ -dimensional framefield which effectively reduces it to a d -dimensional spatial framefield $e_{(i)j}$.

During the following calculations we neglect the π_{matter}^{ij} terms, which are far too high in their order, such that the modified source terms are given by [76, Eqs. (6.33) and (6.34)]

$$\mathcal{H}^{\text{matter}} = \sum_a \left[-n\hat{p}_a \delta_a + \frac{\hat{p}_{aj} \gamma^{ji}}{n\hat{p}_a} \hat{A}_a^{k\ell} e_{(m)k} e_{\ell,i}^{(m)} \delta_a \right. \\ \left. + \left\{ \frac{1}{2} \left(\frac{e_{(r)\ell} e_{(s)i} \hat{p}_{aj}}{n\hat{p}_a} + \gamma^{mn} \frac{e_{(r)m} e_{(s)i} \hat{p}_{aj} \hat{p}_{an} \hat{p}_{a\ell}}{(n\hat{p}_a)^2 (m_a - n\hat{p}_a)} \right) \gamma^{k\ell} \left(\gamma^{nj(d)} \Gamma_{nk}^i + \gamma^{in(d)} \Gamma_{nk}^j \right) \delta_a \right. \right. \\ \left. \left. - \left(\frac{\hat{p}_{a\ell}}{m_a - n\hat{p}_a} \gamma^{ij} \gamma^{k\ell} e_{(r)j} e_{(s)k} \delta_a \right)_{,i} \right\} \hat{S}_{a(r)(s)} \right], \quad (94)$$

$$\mathcal{H}_i^{\text{matter}} = \sum_a \left[\hat{p}_{ai} \delta_a - \hat{A}_a^{k\ell} e_{(m)k} e_{\ell,i}^{(m)} \delta_a + \frac{1}{2} \left(\gamma^{mk} e_{(r)i} e_{(s)k} \delta_a \right. \right.$$

⁴Such frames were originally invented by Élie Cartan and named “repère mobile”, which is French for “moving frame”. In $d = 3$ it is also called “triad” or “dreibein”; in $d = 4$ “tetrad” or “vierbein”. For arbitrary integer d it is called “vielbein”.

$$- \frac{\hat{p}_a \ell \hat{p}_a k}{n \hat{p}_a (m_a - n \hat{p}_a)} (\gamma^{mk} \delta_i^p + \gamma^{mp} \delta_i^k) \gamma^{q\ell} e_{(r)q} e_{(s)p} \delta_a \Big]_{,m} \hat{S}_{a(r)(s)} \Big], \quad (95)$$

where m_a is the mass of the a th object, $\hat{\mathbf{p}}_a$ its canonical momentum, $\hat{S}_{a(r)(s)}$ its canonical spin, $n \hat{p}_a = -\sqrt{m_a^2 + \gamma^{ij} \hat{p}_{a i} \hat{p}_{a j}}$, and $\delta_a = \delta(\mathbf{x} - \hat{\mathbf{z}}_a)$ is the d -dimensional Dirac delta located at $\mathbf{x} = \hat{\mathbf{z}}_a$. Furthermore

$$\hat{A}_a^{k\ell} = \gamma^{ik} \gamma^{j\ell} \left(\frac{1}{2} \hat{S}_{a ij} + \frac{m_a \hat{p}_a (i(n S_a)_j)}{n \hat{p}_a (m_a - n \hat{p}_a)} \right), \quad (96)$$

$$(n S_a)_i = - \frac{\hat{p}_{a j} \gamma^{jk} \hat{S}_{a ki}}{m_a}, \quad (97)$$

to linear order in spin. However the $\hat{A}_a^{k\ell}$ -terms do not contribute to the expanded source expressions at the orders considered here. The matter position and matter momentum variables are canonical conjugate to each other, namely

$$\{\hat{z}_a^i, \hat{p}_{b j}\} = \delta_{ij} \delta_{ab}, \quad (98)$$

and the spin variables fulfill also canonical Poisson bracket relations, namely

$$\{\hat{S}_{a(i)(j)}, \hat{S}_{a(k)(\ell)}\} = \delta_{ik} \hat{S}_{a(j)(\ell)} - \delta_{i\ell} \hat{S}_{a(j)(k)} - \delta_{jk} \hat{S}_{a(i)(\ell)} + \delta_{j\ell} \hat{S}_{a(i)(k)}, \quad (99)$$

where the canonical spin tensor $\hat{S}_{a(i)(j)}$ is related to the canonical spin vector $\hat{\mathbf{S}}_a$ via $\hat{S}_{a(i)(j)} = \varepsilon_{ijk} \hat{S}_a^{(k)}$ and ε_{ijk} is the Levi-Civita symbol. The appropriate Poisson brackets for the canonical spin vector are given by

$$\{\hat{S}_a^{(i)}, \hat{S}_a^{(j)}\} = \varepsilon_{ijk} \hat{S}_a^{(k)}. \quad (100)$$

5.1 Framefield Expansion

We choose to work within a symmetrical framefield gauge $e_{(i)j} = e_{(j)i}$ [188], so the dreibein can be written as a matrix square root of the metric, symbolically $e_{(i)j} = \sqrt{\gamma_{ij}}$ or more explicit

$$e_{(i)k} e_{(k)j} = \gamma_{ij}. \quad (101)$$

Notice that γ_{ij} is positive definite and we require the same for $e_{(i)j}$ such that it is unique. The second relation, (101), can be inverted order by order, namely

$$e_{(0)(i)k} e_{(0)(k)j} = \gamma_{(0)ij} \xrightarrow{\text{ADMTT}} e_{(0)(i)j} = \delta_{ij}, \quad (102a)$$

$$e_{(2)(i)k} e_{(0)(k)j} + e_{(0)(i)k} e_{(2)(k)j} = \gamma_{(2)ij} \xrightarrow{\text{ADMTT}} e_{(2)(i)j} = \frac{1}{2} \gamma_{(2)ij} = \frac{2}{d-2} \bar{\phi}_{(2)} \delta_{ij}, \quad (102b)$$

:

and at the end of the day one gets

$$e_{(0)(i)j} = \delta_{ij}, \quad (103a)$$

$$e_{(2)(i)j} = \frac{2}{d-2} \bar{\phi}_{(2)} \delta_{ij}, \quad (103b)$$

$$e_{(4)(i)j} = \left(\frac{2}{d-2} \bar{\phi}_{(4)} - \frac{d-4}{(d-2)^2} \bar{\phi}_{(2)}^2 \right) \delta_{ij} + \frac{1}{2} h_{ij}^{\text{TT}}, \quad (103c)$$

$$e_{(6)(i)j} = \left(\frac{2}{d-2} \bar{\phi}_{(6)} - \frac{d-4}{(d-2)^2} \bar{\phi}_{(2)} \bar{\phi}_{(4)} + \frac{2}{3} \frac{(d-4)(d-3)}{(d-2)^3} \bar{\phi}_{(2)}^3 \right) \delta_{ij} - \frac{1}{d-2} \bar{\phi}_{(2)} h_{ij}^{\text{TT}}, \quad (103d)$$

for the framefield perturbations. The antisymmetric part of the framefield (which is zero in this gauge) can be interpreted as rotational degrees of freedom in the choice of the local frame. Such a rotation does not change the length of the spins. Recall that an antisymmetric matrix is an infinitesimal generator of rotations and in d dimensions has $\frac{1}{2} d(d-1)$ independent entries. This is exactly the number of rotation planes in d dimensions.

5.2 Constraint Sources

After performing the expansion of the sources (94) and (95) in powers of c^{-1} and also expanding the metric-framefield relation (101), we are able to write down the appropriate post-Newtonian contributions to the source of the Hamilton constraint, namely

$$\mathcal{H}_{(2)}^{\text{matter}} = \sum_a m_a \delta_a, \quad \mathcal{H}_{(4)}^{\text{matter}} = \sum_a \left[\frac{\hat{\mathbf{p}}_a^2}{2m_a} \delta_a + \frac{1}{2m_a} \hat{p}_{ai} \hat{S}_{a(i)(j)} \delta_{a,j} \right], \quad (104a)$$

$$\mathcal{H}_{(6)}^{\text{matter}} = \sum_a \left[-\frac{(\hat{\mathbf{p}}_a^2)^2}{8m_a^3} \delta_a - \frac{2}{d-2} \frac{\hat{\mathbf{p}}_a^2}{m_a} \bar{\phi}_{(2)} \delta_a + \frac{2}{d-2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(j)} \bar{\phi}_{(2),j} \delta_a - \frac{\hat{\mathbf{p}}_a^2}{8m_a^3} \hat{p}_{ai} \hat{S}_{a(i)(j)} \delta_{a,j} \right. \\ \left. - \frac{2}{d-2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(j)} (\bar{\phi}_{(2)} \delta_a)_{,j} \right], \quad (104b)$$

$$\mathcal{H}_{(8)}^{\text{matter}} = \sum_a \left[\frac{(\hat{\mathbf{p}}_a^2)^3}{16m_a^5} \delta_a + \frac{1}{d-2} \frac{(\hat{\mathbf{p}}_a^2)^2}{m_a^3} \bar{\phi}_{(2)} \delta_a + \frac{d+2}{d-2} \frac{\hat{\mathbf{p}}_a^2}{m_a} \bar{\phi}_{(2)}^2 \delta_a - \frac{2}{d-2} \frac{\hat{\mathbf{p}}_a^2}{m_a} \bar{\phi}_{(4)} \delta_a \right. \\ \left. - \frac{1}{2m_a} \hat{p}_{ai} \hat{p}_{aj} h_{ij}^{\text{TT}} \delta_a - \frac{1}{d-2} \frac{\hat{\mathbf{p}}_a^2}{m_a^3} \hat{p}_{ai} \hat{S}_{a(i)(j)} \bar{\phi}_{(2),j} \delta_a - \frac{2(d+2)}{(d-2)^2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(j)} \bar{\phi}_{(2)} \bar{\phi}_{(2),j} \delta_a \right. \\ \left. + \frac{2}{d-2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(j)} \bar{\phi}_{(4),j} \delta_a + \frac{1}{2m_a} \hat{p}_{ai} \hat{S}_{a(j)(k)} h_{ij,k}^{\text{TT}} \delta_a \right] + \sum_a \partial_j \left[\frac{(\hat{\mathbf{p}}_a^2)^2}{16m_a^5} \hat{p}_{ai} \hat{S}_{a(i)(j)} \delta_a \right. \\ \left. + \frac{1}{d-2} \frac{\hat{\mathbf{p}}_a^2}{m_a^3} \hat{p}_{ai} \hat{S}_{a(i)(j)} \bar{\phi}_{(2)} \delta_a + \frac{d+2}{(d-2)^2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(j)} \bar{\phi}_{(2)}^2 \delta_a - \frac{2}{d-2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(j)} \bar{\phi}_{(4)} \delta_a \right. \\ \left. + \frac{1}{4m_a} \hat{p}_{ai} \hat{S}_{a(k)(i)} h_{jk}^{\text{TT}} \delta_a - \frac{1}{4m_a} \hat{p}_{ai} \hat{S}_{a(k)(j)} h_{ik}^{\text{TT}} \delta_a \right], \quad (104c)$$

$$\mathcal{H}_{(10)}^{\text{matter}} = \sum_a \left[-\frac{5(\hat{\mathbf{p}}_a^2)^4}{128m_a^7} \delta_a - \frac{3}{4(d-2)} \frac{(\hat{\mathbf{p}}_a^2)^3}{m_a^5} \bar{\phi}_{(2)} \delta_a - \frac{d+6}{2(d-2)^2} \frac{(\hat{\mathbf{p}}_a^2)^2}{m_a^3} \bar{\phi}_{(2)}^2 \delta_a \right. \\ \left. - \frac{2d(d+2)}{3(d-2)^3} \frac{\hat{\mathbf{p}}_a^2}{m_a} \bar{\phi}_{(2)}^3 \delta_a + \frac{1}{d-2} \frac{(\hat{\mathbf{p}}_a^2)^2}{m_a^3} \bar{\phi}_{(4)} \delta_a + \frac{2(d+2)}{(d-2)^2} \frac{\hat{\mathbf{p}}_a^2}{m_a} \bar{\phi}_{(2)} \bar{\phi}_{(4)} \delta_a \right. \\ \left. - \frac{2}{d-2} \frac{\hat{\mathbf{p}}_a^2}{m_a} \bar{\phi}_{(6)} \delta_a + \frac{\hat{\mathbf{p}}_a^2}{4m_a^3} \hat{p}_{ai} \hat{p}_{aj} h_{ij}^{\text{TT}} \delta_a + \frac{4}{d-2} \frac{\hat{p}_{ai} \hat{p}_{aj}}{m_a} h_{ij}^{\text{TT}} \bar{\phi}_{(2)} \delta_a \right. \\ \left. + \frac{3}{4(d-2)} \frac{(\hat{\mathbf{p}}_a^2)^2}{m_a^5} \hat{p}_{ai} \hat{S}_{a(i)(j)} \bar{\phi}_{(2),j} \delta_a + \frac{d+6}{(d-2)^2} \frac{\hat{\mathbf{p}}_a^2}{m_a^3} \hat{p}_{ai} \hat{S}_{a(i)(j)} \bar{\phi}_{(2)} \bar{\phi}_{(2),j} \delta_a \right. \\ \left. + \frac{2d(d+2)}{(d-2)^3} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(j)} \bar{\phi}_{(2)}^2 \bar{\phi}_{(2),j} \delta_a - \frac{1}{d-2} \frac{\hat{\mathbf{p}}_a^2}{m_a^3} \hat{p}_{ai} \hat{S}_{a(i)(j)} \bar{\phi}_{(4),j} \delta_a \right. \\ \left. - \frac{2(d+2)}{(d-2)^2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(j)} (\bar{\phi}_{(2)} \bar{\phi}_{(4)})_{,j} \delta_a + \frac{2}{d-2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(j)} \bar{\phi}_{(6),j} \delta_a \right. \\ \left. - \frac{\hat{\mathbf{p}}_a^2}{4m_a^3} \hat{p}_{ai} \hat{S}_{a(j)(k)} h_{ij,k}^{\text{TT}} \delta_a - \frac{4}{d-2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(j)(k)} h_{ij,k}^{\text{TT}} \bar{\phi}_{(2)} \delta_a \right. \\ \left. + \frac{3}{d-2} \frac{\hat{p}_{ai}}{m_a} h_{ik}^{\text{TT}} \hat{S}_{a(j)(k)} \bar{\phi}_{(2),j} \delta_a - \frac{1}{d-2} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(i)(k)} h_{jk}^{\text{TT}} \bar{\phi}_{(2),j} \delta_a \right] + (\text{td}). \quad (104d)$$

In $\mathcal{H}_{(10)}^{\text{matter}}$ there is a $\bar{\phi}_{(6)}$ term left. But all occurrences of $\bar{\phi}_{(6)}$ can be cast into the form $-\frac{4}{d-2} \mathcal{H}_{(4)}^{\text{matter}} \bar{\phi}_{(6)}$ which we integrate by parts using (47). Then $\bar{\phi}_{(6)}$ disappears and gets substituted by

$$-\frac{4}{d-2} \mathcal{H}_{(4)}^{\text{matter}} \bar{\phi}_{(6)} = -\frac{4}{d-2} \left[\sum_a \left\{ \frac{(\hat{\mathbf{p}}_a^2)^2}{16m_a^3} \bar{S}_{(4)} \delta_a + \frac{d+2}{4(d-2)} \frac{\hat{\mathbf{p}}_a^2}{m_a} \bar{\phi}_{(2)} \bar{S}_{(4)} \delta_a \right. \right. \\ \left. \left. + \frac{m_a}{2} (\bar{\phi}_{(4)} - \bar{\phi}_{(2)}^2) \bar{S}_{(4)} \delta_a + \frac{\hat{\mathbf{p}}_a^2 \hat{p}_{ai} \hat{S}_{a(i)(j)}}{16m_a^3} \bar{S}_{(4)} \delta_{a,j} \right\} \right]$$

$$\begin{aligned}
& + \frac{d+2}{4(d-2)} \frac{\hat{p}_{ai} \hat{S}_{a(i)(j)}}{m_a} \bar{\phi}_{(2)} \bar{S}_{(4)} \delta_{a,j} \Big\} \\
& - \frac{1}{16\pi G^{(d)}} \left\{ \frac{1}{2} (\tilde{\pi}_{(3)}^{ij})^2 \bar{S}_{(4)} - 2h_{ij}^{\text{TT}} \bar{S}_{(4)} \bar{\phi}_{(2),ij} \right\} \Big] + (\text{td}) .
\end{aligned} \tag{105}$$

Furthermore we are also able to write down the sources for the momentum constraint in their full form, which are given by

$$\mathcal{H}_{i(3)}^{\text{matter}} = \sum_a \left[\hat{p}_{ai} \delta_a + \frac{1}{2} (\hat{S}_{a(i)(j)} \delta_a)_{,j} \right], \tag{106a}$$

$$\mathcal{H}_{i(5)}^{\text{matter}} = \frac{1}{2} \sum_a \left[-\frac{\hat{p}_{ak}}{2m_a^2} (\hat{p}_{aj} \hat{S}_{a(i)(k)} + \hat{p}_{ai} \hat{S}_{a(j)(k)}) \delta_a \right]_{,j}, \tag{106b}$$

$$\begin{aligned}
\mathcal{H}_{i(7)}^{\text{matter}} = & \frac{1}{2} \sum_a \left[\left(-\frac{1}{2} (h_{jk}^{\text{TT}} \hat{S}_{a(i)(k)} + h_{ik}^{\text{TT}} \hat{S}_{a(j)(k)}) + \frac{3\hat{\mathbf{p}}_a^2}{8m_a^4} \hat{p}_{ak} (\hat{p}_{aj} \hat{S}_{a(i)(k)} + \hat{p}_{ai} \hat{S}_{a(j)(k)}) \right. \right. \\
& \left. \left. + \frac{2}{d-2} \frac{\hat{p}_{ak}}{m_a^2} (\hat{p}_{aj} \hat{S}_{a(i)(k)} + \hat{p}_{ai} \hat{S}_{a(j)(k)}) \bar{\phi}_{(2)} \right) \delta_a \right]_{,j}.
\end{aligned} \tag{106c}$$

With the source expressions (104) including (105) and (106) from above, we may express $B_{(4)ij}$ and $\hat{B}_{(6)ij}$ in terms of the matter variables. They are given by

$$B_{(4)ij} = 16\pi G^{(d)} \sum_a \left[-\frac{\hat{p}_{ai} \hat{p}_{aj}}{2m_a} \delta_a - \frac{\hat{p}_{ai} \hat{S}_{a(j)(k)}}{2m_a} \delta_{a,k} \right] - \frac{4(d-1)}{d-2} \bar{\phi}_{(2),i} \bar{\phi}_{(2),j}, \tag{107a}$$

$$\begin{aligned}
\hat{B}_{(6)ij} = & 16\pi G^{(d)} \sum_a \left[\frac{\hat{\mathbf{p}}_a^2}{4m_a^3} \hat{p}_{ai} \hat{p}_{aj} \delta_a + \frac{d+2}{d-2} \frac{\hat{p}_{ai} \hat{p}_{aj}}{m_a} \bar{\phi}_{(2)} \delta_a \right. \\
& + \frac{\hat{\mathbf{p}}_a^2}{4m_a^3} \hat{p}_{ai} \hat{S}_{a(j)(k)} \delta_{a,k} + \frac{d+2}{d-2} \frac{\hat{p}_{ai} \hat{S}_{a(j)(k)}}{m_a} (\bar{\phi}_{(2)} \delta_a)_{,k} \\
& + \frac{d+4}{2(d-2)} \frac{\hat{p}_{ai}}{m_a} \hat{S}_{a(k)(j)} \bar{\phi}_{(2),k} \delta_a - \frac{d}{2(d-2)} \frac{\hat{p}_{ak}}{m_a} \hat{S}_{a(k)(j)} \bar{\phi}_{(2),i} \delta_a \\
& \left. + \frac{1}{2} (V_{(3),k}^j + V_{(3),j}^k) \hat{S}_{a(k)(i)} \delta_a \right] \\
& + 2\tilde{\pi}_{(3)}^{ik} (\tilde{\pi}_{(3),j}^k - \tilde{\pi}_{(3),k}^j) + \frac{d-2}{d-1} \tilde{\pi}_{(3)}^{ij} \tilde{\pi}_{(3),k}^k + 2\tilde{\pi}_{(3),k}^{ij} V_{(3)}^k \\
& + 8\frac{2d-3}{d-2} \bar{\phi}_{(4)} \bar{\phi}_{(2),ij} + 8\frac{3d-4}{(d-2)^2} \bar{\phi}_{(2)} \bar{\phi}_{(2),i} \bar{\phi}_{(2),j} + 4\frac{d-4}{d-2} \bar{S}_{(4)} \bar{\phi}_{(2),ij}.
\end{aligned} \tag{107b}$$

We expressed $\hat{B}_{(6)ij}$ in a more convenient way now, since the $\tilde{\pi}_{(3)}^i$ vector potential has a much more simple structure than the $V_{(3)}^i$ vector potential.

5.3 Matter Correction to the Canonical Field Momentum

Since we eliminated the transverse-traceless part of the canonical field momentum via using the relation between canonical field momentum and velocity of the h_{ij}^{TT} field (71), there are terms containing matter parts of the field momentum left in the Routhian (93b). These can be calculated from e.g. [77, Eq. (2.34)] where π_{matter}^{ij} is given by

$$\pi_{\text{matter}}^{ij} = 16\pi G^{(d)} \sum_a \pi_a^{ij} \delta_a. \tag{108}$$

From [77, Eqs. (3.33) and (3.34)] one gets the closed form expression

$$\pi_a^{ij} = \frac{1}{2} \gamma^{ik} \gamma^{j\ell} \frac{m_a \hat{p}_{a(k} n S_{a\ell)}}{n \hat{p}_a (m_a - n \hat{p}_a)}. \tag{109}$$

The part containing the antisymmetric $\hat{A}_a^{[ij]}$ was neglected, because it is of order c^{-7} ($B_{k\ell}^{ij}$ starts at c^{-4}) which is not necessary here. Here we only need π_a^{ij} to the order c^{-5} . A power counting (see beginning of subsection 3.1) tells us that we only have to take the leading order approximation of the above expression, reading

$$\pi_{(5)a}^{ij} = \frac{\hat{p}_{a\ k}}{8m_a^2} \left(\hat{p}_{a\ i} \hat{S}_{a\ (k)(j)} + \hat{p}_{a\ j} \hat{S}_{a\ (k)(i)} \right). \quad (110)$$

This expression has a vanishing trace, which makes obvious that we can neglect $\tilde{\pi}^{ij}$, Eq. (32), at the considered order.

6 Field Solutions and Integration

After obtaining expressions for the sources (104), (106) and the Routhian (93) which gives the Hamiltonian in the matter degrees of freedom after an integration, we need the fields to be inserted into the Routhian. These can be derived by solving the lower order constraint equations in case of the post-Newtonian potential and the non-propagating parts of the field momentum (see 6.1). For the propagating degrees of freedom the wave equation has to be solved (see 6.2).

6.1 d -dimensional Solutions of the Constraints

With $K = \frac{\Gamma(\frac{d}{2}-1)}{\pi^{\frac{d}{2}-1}} G^{(d)}$, the Hamilton constraint equations (43a), (43b), the momentum constraint equation (45a), and the various transformation formulas (33), (34), and (35) relating the longitudinal field momentum and its corresponding vector potentials, we find using the inverse Laplacians listed in the Appendix A.1 that

$$\phi_{(2)} = 4K \sum_a \frac{m_a}{r_a^{d-2}}, \quad (111)$$

$$\phi_{(4)} = 4K \sum_a \left[\frac{\hat{\mathbf{p}}_a^2}{2m_a} \frac{1}{r_a^{d-2}} + \frac{\hat{p}_{a\ i} \hat{S}_{a\ (i)(j)}}{2m_a} \left(\frac{1}{r_a^{d-2}} \right)_{,j} - K \frac{d-2}{d-1} \sum_{b \neq a} \frac{m_a m_b}{r_{ab}^{d-2} r_b^{d-2}} \right], \quad (112)$$

$$\tilde{\pi}_{(3)}^i = K \sum_a \left[2 \frac{\hat{p}_{a\ i}}{r_a^{d-2}} + \hat{S}_{a\ (i)(j)} \left(\frac{1}{r_a^{d-2}} \right)_{,j} \right], \quad (113)$$

$$V_{(3)}^i = K \sum_a \left[2 \frac{\hat{p}_{a\ i}}{r_a^{d-2}} - \frac{d-2}{2(d-1)(4-d)} \hat{p}_{a\ j} \left(\frac{1}{r_a^{d-4}} \right)_{,ij} + \hat{S}_{a\ (i)(j)} \left(\frac{1}{r_a^{d-2}} \right)_{,j} \right], \quad (114)$$

$$\begin{aligned} \tilde{\pi}_{(3)}^{ij} = K \sum_a \left[2 \hat{p}_{a\ i} \left(\frac{1}{r_a^{d-2}} \right)_{,j} + 2 \hat{p}_{a\ j} \left(\frac{1}{r_a^{d-2}} \right)_{,i} - \frac{d-2}{(d-1)(4-d)} \hat{p}_{a\ k} \left(\frac{1}{r_a^{d-4}} \right)_{,ijk} \right. \\ \left. - \frac{2}{d-1} \delta_{ij} \hat{p}_{a\ k} \left(\frac{1}{r_a^{d-2}} \right)_{,k} + \hat{S}_{a\ (i)(k)} \left(\frac{1}{r_a^{d-2}} \right)_{,kj} + \hat{S}_{a\ (j)(k)} \left(\frac{1}{r_a^{d-2}} \right)_{,ki} \right]. \end{aligned} \quad (115)$$

Remember that the momentum constraint can be solved for $\tilde{\pi}^i$ with the help of (36). The more complicated fields like $\tilde{\pi}_{(5)}^i$ or $\phi_{(6)}$ were so far only found in $d = 3$ dimensions [27]. Also the leading order of the transverse-traceless part of the metric is only partially known in d dimensions. We will discuss these issues in the following subsection.

6.2 Solutions of the Wave Equation

Consider now the wave equation (70) for h_{ij}^{TT} . There h_{ij}^{TT} is given in terms of a post-Newtonian approximate source S_{ij} , namely

$$\square h_{ij}^{\text{TT}} = \delta_{k\ell}^{\text{TT}ij} (S_{(4)k\ell} + S_{(6)k\ell}). \quad (116)$$

By taking into account the near-zone expansion of h_{ij}^{TT} , (79), one gets

$$h_{(4)ij}^{\text{TT}} = \delta_{k\ell}^{\text{TT}ij} \Delta^{-1} S_{(4)k\ell}, \quad (117)$$

$$h_{(6)ij}^{\text{TT}} = \delta_{k\ell}^{\text{TT}ij} \Delta^{-1} (S_{(6)k\ell} + \Delta^{-1} \partial_t^2 S_{(4)k\ell}) = \Delta^{-1} \left(\delta_{k\ell}^{\text{TT}ij} S_{(6)k\ell} + \ddot{h}_{(4)ij}^{\text{TT}} \right), \quad (118)$$

for the leading order and next-to-leading order expressions of h_{ij}^{TT} in the near-zone. Here the sources are given by

$$S_{(4)ij} = 2B_{(4)ij}, \quad (119)$$

$$S_{(6)ij} = 2\hat{B}_{(6)ij} - \frac{16\pi G^{(d)}}{d-1} \mathcal{H}_{(2)}^{\text{matter}} h_{(4)ij}^{\text{TT}} + \frac{8}{d-2} (\bar{\phi}_{(2)} h_{(4)ij,k}^{\text{TT}})_{,k} - \frac{\partial}{\partial t} C_{(5)ij}, \quad (120)$$

where $B_{(4)ij}$ and $\hat{B}_{(6)ij}$ are given by (107a) and (107b), and $C_{(5)ij}$ by (65) via (108) and (110). Notice that we removed the post-Newtonian order-counting parameter c in (117) and (118). Fortunately there is no need to evaluate (118) here. In fact the $h_{(4)ij}^{\text{TT}}$ dependence of (120) renders the calculation almost impossible.

Then the solution of the wave equation at leading order and linear in spin is given by

$$\begin{aligned} h_{(4)ij}^{\text{TT}} = & 4K \sum_a \left[\frac{\hat{p}_a i \hat{p}_a j}{m_a} \frac{1}{r_a^{d-2}} - \frac{1}{4-d} \frac{\hat{p}_a k \hat{p}_a (i}{m_a} \left(\frac{1}{r_a^{d-4}} \right)_{,j)k} \right. \\ & - \frac{1}{d-1} \delta_{ij} \left(\frac{\hat{\mathbf{p}}_a^2}{m_a} \frac{1}{r_a^{d-2}} - \frac{1}{2(4-d)} \frac{\hat{p}_a k \hat{p}_a \ell}{m_a} \left(\frac{1}{r_a^{d-4}} \right)_{,k\ell} \right) \\ & + \frac{1}{2(d-1)(4-d)} \frac{\hat{\mathbf{p}}_a^2}{m_a} \left(\frac{1}{r_a^{d-4}} \right)_{,ij} \\ & + \frac{d-2}{8(d-1)(d-4)(d-6)} \frac{\hat{p}_a k \hat{p}_a \ell}{m_a} \left(\frac{1}{r_a^{d-6}} \right)_{,ijk\ell} \\ & + \frac{\hat{p}_a k \hat{S}_a(\ell)(m)}{m_a} \left\{ \left(\delta_{k(i} \delta_{j)\ell} \partial_m - \frac{1}{d-1} \delta_{ij} \delta_{k\ell} \partial_m \right) \frac{1}{r_a^{d-2}} \right. \\ & \left. \left. + \frac{1}{2(4-d)} \left(\frac{1}{d-1} \delta_{k\ell} \partial_i \partial_j \partial_m - \delta_{\ell(i} \partial_{j)} \partial_k \partial_m \right) \frac{1}{r_a^{d-4}} \right\} \right] + h_{(40)ij}^{\text{TT}}, \end{aligned} \quad (121)$$

where $h_{(40)ij}^{\text{TT}}$ is the momentum (and spin-) independent part of the transverse-traceless part of the metric which is generated by the TT-projection of $\Delta^{-1}(\bar{\phi}_{(2),i} \bar{\phi}_{(2),j})$ and is only known in $d=3$ see [27, Eq. (A20)], namely

$$\begin{aligned} h_{(40)ij}^{\text{TT}} = & G^2 \sum_a \sum_{b \neq a} m_a m_b \left\{ -\frac{4}{s_{ab}} \left(\frac{1}{r_{ab}} + \frac{1}{s_{ab}} \right) n_{ab}^i n_{ab}^j + \frac{1}{4} \left(\frac{r_a + r_b}{r_{ab}^3} + \frac{12}{s_{ab}^2} \right) n_a^i n_b^j \right. \\ & + 2 \left(\frac{2}{s_{ab}^2} - \frac{1}{r_{ab}^2} \right) (n_a^i n_{ab}^j + n_a^j n_{ab}^i) \\ & + \left[\frac{5}{8r_{ab}r_a} - \frac{1}{8r_{ab}^3} \left(\frac{r_b^2}{r_a} + 3r_a \right) - \frac{1}{s_{ab}} \left(\frac{1}{r_a} + \frac{1}{s_{ab}} \right) \right] n_a^i n_a^j \\ & \left. + \left[\frac{5r_a}{8r_{ab}^3} \left(\frac{r_a}{r_b} - 1 \right) - \frac{17}{8r_{ab}r_a} + \frac{1}{2r_a r_b} + \frac{1}{s_{ab}} \left(\frac{1}{r_a} + \frac{4}{r_{ab}} \right) \right] \delta_{ij} \right\}. \end{aligned} \quad (122)$$

Both solutions were also obtained by using the inverse Laplacians in the Appendix A.1. Obviously, most of the parts of $h_{(6)ij}^{\text{TT}}$ are of the same type as $h_{(40)ij}^{\text{TT}}$ (see (118) and (120)). That is the reason why we eliminated it from the integrands.

6.3 Distributional Contributions

As long as the Riesz-kernel method is not used (where a Dirac delta is substituted by the so-called Riesz-kernel) one has to take care of delta parts when differentiating certain functions. Consider for example the

field $\phi_{(2)} = 4K \sum_a \frac{m_a}{r_a^{d-2}}$ and differentiate it two times. Using the ordinary derivative it would give

$$\partial_i^{\text{ord}} \partial_j^{\text{ord}} \phi_{(2)} = 0. \quad (123)$$

But as we already know from the constraint equations the second derivative of $\phi_{(2)}$ should be

$$\partial_i \partial_j \phi_{(2)} = -16\pi G^{(d)} \frac{1}{d} \delta_{ij} \sum_a m_a \delta_a. \quad (124)$$

Fortunately there is a result from the theory of distributions [27] which defines the so-called distributional derivative

$$\partial_i f = \partial_i^{\text{ord}} f + \frac{(-1)^k}{k!} \frac{\partial^k \delta(\mathbf{x})}{\partial x^{i_1} \dots \partial x^{i_k}} \oint_{\Sigma} d\Omega_{d-1} n^i f x^{i_1} \dots x^{i_k}. \quad (125)$$

Here f is a positive homogeneous function of degree λ (i.e. $f(a\mathbf{x}) = a^\lambda f(\mathbf{x})$ for $a \geq 0$) and $k := -\lambda + 1 - d$ is a non-negative integer. This means f must decay with an exponent linear in the dimension d which does not apply to fields generated by a Riesz-kernel type source (see Appendix A.4). There are not only distributional contributions from the field derivatives, but from the fields themselves (some parts of the higher order field momenta).

6.4 Ultraviolet-Analysis

As for gauge theories in quantum field theory, dimensional regularization should be used in classical general relativity [31]. Therefore first all integrals must be evaluated in generic d dimensions and then the limit $d \rightarrow 3$ is calculated. However, certain integrals are very difficult to solve for generic d . In practice one therefore evaluates the integrals in $d = 3$ first and then determines possible additional contributions that arise from dimensional regularization. That is, one analyses the d -dependence of the integrals close to the singular sources, i.e., in the UV. (Only close to singularities regularization is important.) This is the purpose of the present section. Other necessary integration techniques are provided in Appendix A.

The UV-analysis in generic dimension d is a necessary ingredient to correctly derive the Hamiltonians at formal 3PN level. This includes the 3PN point-mass Hamiltonian (see [31]) and the NNLO spin-orbit and spin(1)-spin(2) Hamiltonians considered in the present article. It would also be necessary for the yet unknown NNLO spin(1)-spin(1) Hamiltonian.

For integrals only obtained for $d = 3$ one has no control on poles in $1/(d - 3)$. There are two different problems with such poles: First the poles do not appear in pure $d = 3$ calculations and thus lead to ambiguous results after integrations by parts in integrands containing such poles (in one representation there are poles, in another maybe not). This comes from the fact that some of the pole terms can also give finite contributions which must be added to the $d = 3$ result. Second the poles have to cancel each other in order to extract a finite result from the d dimensional integration in the limit $d \rightarrow 3$ (or one must be able to absorb all poles through a renormalization procedure as in [184]). Both problems are well-known and also discussed in [31]. In the following we will provide some more technical details on how to perform the UV-analysis depending on the structure of the integrand.

All integrals involving $h_{(40)}^{\text{TT}}{}_{ij}$, $\delta_{kl}^{\text{TT}ij}(\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{kl})$ and the high order potentials such as $\bar{\phi}_{(6)}$ or $\tilde{\pi}_{(5)}^i$ are not available in d dimensions and were only calculated in $d = 3$ dimensions here. In all other integrals the limit $d \rightarrow 3$ is straightforward, although integrations in d dimensions sometimes involve around one million terms on which the limit must be performed. In case of the TT-projection of $\bar{\phi}_{(2)} \tilde{\pi}_{(3)}^{ij}$, the fields are available in d dimensions. Hence, one can split up this part of the Hamiltonian in one-particle TT-projections (which can be performed in d dimensions) and two-particle TT-projections (which can only be evaluated for $d = 3$ with the presented methods). For the latter ones must still perform the UV-analysis.

The term UV-analysis in this context refers to the short-range behavior of the integrand around a specific point. This will become more clear during the following explanation. Let us now consider the decay of the integrand $f(r_a, r_b, \mathbf{n}_a, \mathbf{n}_b)$ around the source a . First of all the integral is split up into a ball integral around

one of the sources, say for example source particle a , and an integral over the whole $\mathbb{R}^d$ without this ball,

$$\int d^d x f(r_a, r_b, \mathbf{n}_a, \mathbf{n}_b) = \int_{B_{\ell_a}(\hat{\mathbf{z}}_a)} d^d x f(r_a, r_b, \mathbf{n}_a, \mathbf{n}_b) + \int_{\mathbb{R}^d \setminus B_{\ell_a}(\hat{\mathbf{z}}_a)} d^d x f(r_a, r_b, \mathbf{n}_a, \mathbf{n}_b), \quad (126)$$

where $0 < \ell_a \ll r_{ab}$. The variables r_b and $\mathbf{n}_b$ of the other source ($b \neq a$) are expressed in terms of r_a , r_{ab} and $\mathbf{n}_{ab}$,

$$r_b = |\mathbf{x} - \hat{\mathbf{z}}_b| = |\mathbf{x} - \hat{\mathbf{z}}_a + \hat{\mathbf{z}}_a - \hat{\mathbf{z}}_b| = \sqrt{r_a^2 + r_{ab}^2 + 2r_a r_{ab}(\mathbf{n}_a \cdot \mathbf{n}_{ab})}, \quad (127)$$

$$\mathbf{n}_b = \frac{r_a}{r_b} \mathbf{n}_a + \frac{r_{ab}}{r_b} \mathbf{n}_{ab}, \quad (128)$$

such that all $\mathbf{x}$ -dependent expressions come from r_a and $\mathbf{n}_a$ type variables. Next we concentrate on the ball integral around a ,

$$\int_{B_{\ell_a}(\mathbf{x}_a)} d^d x f(r_a, \mathbf{n}_a) = \int d\Omega_{a,d-1} \int_0^{\ell_a} dr_a r_a^{d-1} f(r_a, \mathbf{n}_a). \quad (129)$$

Now the integrand is expanded in r_a (leaving $\mathbf{n}_a$ untouched). This is possible because a and b are well separated, and the ball contains only a small neighborhood of the source a . Then the integrand takes the form of a polynomial in r_a and one can pick out the terms contributing poles at $d = 3$ (the ones with an exponent giving -3 for $d = 3$ on r_a). The next step is to count the number of $\mathbf{n}_a$ -vectors in each term and remove terms with an odd number of these vectors. This is due to the averaging procedure coming from the angular integration in (129) using the formulas (172) in the Appendix. Consider for example an integrand $f(r_a, \mathbf{n}_a) = C(d) r_a^{6-3d} (\hat{\mathbf{p}}_a \cdot \mathbf{n}_a) (\hat{\mathbf{p}}_b \cdot \mathbf{n}_a)$ (this integrand is of the form qualified as dangerous in [31, Eq. (3.1)]) then (129) gives

$$\begin{aligned} \int_{B_{\ell_a}(\mathbf{x}_a)} d^d x f(r_a, \mathbf{n}_a) &= C(d) \int d\Omega_{a,d-1} (\hat{\mathbf{p}}_a \cdot \mathbf{n}_a) (\hat{\mathbf{p}}_b \cdot \mathbf{n}_a) \int_0^{\ell_a} dr_a r_a^{d-1} r_a^{6-3d} \\ &= C(d) \hat{p}_{ai} \hat{p}_{bj} \int d\Omega_{a,d-1} n_a^i n_a^j \int_0^{\ell_a} dr_a r_a^{5-2d}. \end{aligned} \quad (130)$$

Using (172b) and usual integration rules we obtain

$$\int_{B_{\ell_a}(\mathbf{x}_a)} d^d x f(r_a, \mathbf{n}_a) = \frac{C(d)}{d} \Omega_{a,d-1} (\hat{\mathbf{p}}_a \cdot \hat{\mathbf{p}}_b) \frac{\ell_a^{2(3-d)}}{2(3-d)}. \quad (131)$$

The last integration step was performed by means of an analytic continuation from $d < 3$. At this stage there are three possibilities: The first one is that $C(d)$ contains several factors of $d - 3$ which cancel the pole in the last factor and even lead to a vanishing limit when $d \rightarrow 3$. Then the potentially dangerous term is actually not dangerous at all. The second possibility is that $C(d) \sim d - 3$ which would also lead to a cancellation of the pole but would give a finite contribution which has to be taken into account to get the correct Hamiltonian. Last but not least $C(d)$ could have such a structure that a pole remains and so this term has to be renormalized or canceled by another pole to give a physically meaningful result.

The procedure mentioned above is only valid if there is no TT-projection (and in particular no $h_{(40)ij}^{\text{TT}}$) appearing. The analysis of the $h_{(40)ij}^{\text{TT}}$ type integrals works as follows (during this discussion we talk about a two-particle system where we consider r_1 as expansion point): The $h_{(40)ij}^{\text{TT}}$ part is given in terms of inverse Laplacians and field variables in d dimensions by

$$h_{(40)ij}^{\text{TT}} = -\frac{8(d-1)}{d-2} \delta_{kl}^{\text{TT}ij} \Delta^{-1}(\bar{\phi}_{(2),k} \bar{\phi}_{(2),l}), \quad (132)$$

see (122) for the explicit solution in $d = 3$. Now one can insert the $\bar{\phi}_{(2)}$ field, given by

$$\bar{\phi}_{(2)} = \frac{d-2}{4(d-1)} (m_1 u_1 + m_2 u_2), \quad (133)$$

where $u_a = -16\pi G^{(d)}\Delta^{-1}\delta_a \sim r_a^{2-d}$. After interchanging TT-projector and inverse Laplacian, one gets $\delta_{kl}^{\text{TT}ij}(u_{a,k}u_{a,l}) = 0$. One can see this by using $u_{a,k}u_{a,l} \sim n_a^k n_a^l r_a^{2-2d}$ which can be rewritten using

$$\partial_i \partial_j r_a^{4-2d} = -2(d-2)(\delta_{ij} - 2(d-1)n_a^i n_a^j) r_a^{2-2d}, \quad (134)$$

as $u_{a,k}u_{a,l} \sim \frac{1}{2(d-1)}(\delta_{ij} r_a^{2-2d} + \frac{1}{2(d-2)}\partial_i \partial_j r_a^{4-2d})$. This will obviously be projected to zero by the TT-projector. Thus, after TT-projection there is only one part left, namely

$$h_{(40)ij}^{\text{TT}} = -2\frac{d-2}{2(d-1)}m_1 m_2 \Delta^{-1} \delta_{kl}^{\text{TT}ij}(u_{1,k}u_{2,l}). \quad (135)$$

Under the TT-projector one can integrate by parts because $\partial_k \delta_{kl}^{\text{TT}ij} = 0$ and obtains

$$h_{(40)ij}^{\text{TT}} = 2\frac{d-2}{2(d-1)}m_1 m_2 \Delta^{-1} \delta_{kl}^{\text{TT}ij}(u_1 \partial_k \partial_l u_2). \quad (136)$$

Notice that the derivative should act on the u_2 term because all quantities will be Taylor expanded in r_1 around $\mathbf{x} = \hat{\mathbf{z}}_1$ and u_1 is already proportional to an r_1 -power. At this stage of processing the $h_{(40)ij}^{\text{TT}}$ terms one is able to use the same analysis procedure as mentioned above: Check whether there are powers of r_1 with exponents smaller than -3 for $d = 3$ and expand $\partial_k \partial_l u_2$ around r_1 to an order sufficient to reach the critical value in the exponent of r_1 . The idea is the expansion of the r_2 variable in the TT-projector such that the Taylor expansion consists only of multiple inverse Laplacians on powers of r_1 , which can be calculated using (168) from the Appendix. Also bear in mind that the inverse Laplacian introduces additional powers of r_1 via the Green's function. The final ball integration can now be performed as discussed above. The same technique can also be used to perform the UV-analysis of terms involving $\delta_{kl}^{\text{TT}ij}(\bar{\phi}_{(2)}\tilde{\pi}_{(3)}^{kl})$.

To check our UV-analysis code we first reproduced the pole coefficients given in [31, Table 1]. The *finite* UV-contribution to the 3PN point-mass Hamiltonian in our representation (93) is given by

$$\begin{aligned} \Delta H_{\text{PM}}^{3\text{PN},\text{UV}}(d) &= \frac{2\Lambda^{6-2d}(d-2)(d+1)(96-40d-28d^2+d^3)\pi^{3-3d/2}\Gamma(\frac{d}{2})^3}{3(d-4)(d-1)^4(d+2)} \times \\ &\quad \frac{(G^{(d)})^3}{r_{12}^d} m_1 m_2 (d(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2 - \hat{\mathbf{p}}_1^2 + d(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2 - \hat{\mathbf{p}}_2^2), \end{aligned} \quad (137)$$

where Λ is an UV-cutoff scale which does not contribute in the limit $d \rightarrow 3$. A similar analysis for the 2PN point-mass Hamiltonian gave no contribution. We found no net contribution to the spin-dependent Hamiltonians, though poles and finite parts appeared in intermediate expressions. That is, Hadamard regularization would have been sufficient to obtain the correct linear-in-spin Hamiltonians presented in the present work. The same situation was also found for the harmonic-gauge calculation of the equations of motion in [85, 86].

7 Results

After discussing the several simplifications given above to reduce the integral of the formal 3PN Routhian to a form which can be handled appropriately, we continue by giving a short description of the integrands showing up at this order. The integrands can be divided into three different types:

- the delta-type $\int d^d x f(\mathbf{x})\delta_1$,
- the Riesz-type $\int d^d x n_1^{i_1} \dots n_1^{i_k} n_2^{j_1} \dots n_2^{j_\ell} r_1^\alpha r_2^\beta$,
- and the generalized Riesz-type $\int d^3 x n_1^{i_1} \dots n_1^{i_k} n_2^{j_1} \dots n_2^{j_\ell} r_1^\alpha r_2^\beta s_{12}^\gamma$.

The solution of these three types of integrals will be shown in Appendix A.

We used our MATHEMATICA code to perform an integration of (93) directly with all fields inserted up to linear order in spin, neglecting $\text{spin}(1)^2$ and $\text{spin}(2)^2$ terms afterwards. From this we obtained the fully

reduced matter Hamiltonians for point masses, for the spin-orbit- and the spin(1)-spin(2)-interaction. There were no $1/(d-3)$ poles for the linear-in-spin Hamiltonians giving rise to finite parts or being singular for $d \rightarrow 3$. Such poles only appeared in some intermediate steps in the UV-analysis (see Sect. 6.4) but finally identically canceled. For the point-mass parts there appeared a finite contribution given in (137). This together with our integration result reproduced the result from the literature exactly.

Now we are able to write down the NNLO linear-in-spin fully reduced Hamiltonians in terms of matter variables only, by using (93) for the Routhian, inserting the sources (104), $\hat{B}_{(6)ij}$ (107b), π_{matter}^{ij} (110), the solution for the constraint fields (111)–(115), (55), (56), and the propagating degrees of freedom (121) and (122). Notice that certain field derivatives may lead to distributional contributions mentioned in 6.3. The integrals can (for a binary) be solved by using the techniques given in Appendix A. Both Hamiltonians are valid for any compact objects like black holes or neutron stars. Their center-of-mass frame versions are given in 7.3 where the gyromagnetic ratios in the spin-orbit case are also given in [132, 133].

7.1 Next-to-next-to-leading Order Spin-Orbit Hamiltonian

The spin-orbit Hamiltonian given in this subsection is the higher order gravitational analogue of the interaction of an electron's spin interacting with the electron's orbital angular momentum in the case of an e.g. hydrogen atom. In quantum electrodynamics this interaction is responsible for the fine structure in the spectrum. Here the spin is obviously no quantum mechanical quantity, it only characterizes the rotation (i.e. its direction and magnitude) of a gravitating mass in the gravitational field of another mass. Notice that the fine structure constant α in electromagnetic theory is substituted by Newton's gravitational constant G here. The result for the NNLO spin-orbit Hamiltonian reads

$$\begin{aligned}
H_{\text{SO}}^{\text{NNLO}} = & \frac{G}{r_{12}^2} \left[\left(\frac{7m_2(\hat{\mathbf{p}}_1^2)^2}{16m_1^5} + \frac{9(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_1^2}{16m_1^4} + \frac{3\hat{\mathbf{p}}_1^2(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{4m_1^3m_2} \right. \right. \\
& + \frac{45(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^3}{16m_1^2m_2^2} + \frac{9\hat{\mathbf{p}}_1^2(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{16m_1^4} - \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{16m_1^2m_2^2} \\
& - \frac{3(\hat{\mathbf{p}}_1^2)(\hat{\mathbf{p}}_2^2)}{16m_1^3m_2} - \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_2^2}{16m_1^2m_2^2} + \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2\hat{\mathbf{p}}_2^2}{4m_1m_2^3} \\
& - \frac{3(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_2^2}{16m_1^2m_2^2} - \frac{3(\hat{\mathbf{p}}_2^2)^2}{16m_1m_2^3} \Big) ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1)\hat{\mathbf{S}}_1) + \left(-\frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_1^2}{2m_1^3m_2} \right. \\
& - \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{4m_1^2m_2^2} + \frac{3\hat{\mathbf{p}}_1^2(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{4m_1^2m_2^2} - \frac{\hat{\mathbf{p}}_1^2(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{2m_1^3m_2} + \frac{(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)^2}{2m_1^2m_2^2} \\
& + \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2\hat{\mathbf{p}}_2^2}{4m_1^2m_2^2} - \frac{(\hat{\mathbf{p}}_1^2)(\hat{\mathbf{p}}_2^2)}{4m_1^2m_2^2} - \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_2^2}{2m_1m_2^3} \\
& - \frac{(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_2^2}{2m_1m_2^3} \Big) ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1) + \left(-\frac{9(\mathbf{n}_{12} \hat{\mathbf{p}}_1)\hat{\mathbf{p}}_1^2}{16m_1^4} + \frac{\hat{\mathbf{p}}_1^2(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{m_1^3m_2} \right. \\
& + \frac{27(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{16m_1^2m_2^2} - \frac{(\mathbf{n}_{12} \hat{\mathbf{p}}_2)(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{8m_1^2m_2^2} - \frac{5(\mathbf{n}_{12} \hat{\mathbf{p}}_1)\hat{\mathbf{p}}_2^2}{16m_1^2m_2^2} \\
& + \frac{(\mathbf{n}_{12} \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_2^2}{m_1m_2^3} \Big) ((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1) \Big] \\
& + \frac{G^2}{r_{12}^3} \left[\left(-\frac{3m_2(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2}{2m_1^2} + \left(-\frac{3m_2}{2m_1^2} + \frac{27m_2^2}{8m_1^3} \right) \hat{\mathbf{p}}_1^2 + \left(\frac{177}{16m_1} + \frac{11}{m_2} \right) (\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2 \right. \right. \\
& + \left(\frac{11}{2m_1} + \frac{9m_2}{2m_1^2} \right) (\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2) + \left(\frac{23}{4m_1} + \frac{9m_2}{2m_1^2} \right) (\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2) \\
& - \left(\frac{159}{16m_1} + \frac{37}{8m_2} \right) \hat{\mathbf{p}}_2^2 \Big) ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1)\hat{\mathbf{S}}_1) + \left(\frac{4(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2}{m_1} + \frac{13\hat{\mathbf{p}}_1^2}{2m_1} \right. \\
& + \frac{5(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{m_2} + \frac{53\hat{\mathbf{p}}_2^2}{8m_2} - \left(\frac{211}{8m_1} + \frac{22}{m_2} \right) (\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)
\end{aligned}$$

$$\begin{aligned}
& - \left(\frac{47}{8m_1} + \frac{5}{m_2} \right) (\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2) \left((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1 \right) + \left(- \left(\frac{8}{m_1} + \frac{9m_2}{2m_1^2} \right) (\mathbf{n}_{12} \hat{\mathbf{p}}_1) \right. \\
& + \left. \left(\frac{59}{4m_1} + \frac{27}{2m_2} \right) (\mathbf{n}_{12} \hat{\mathbf{p}}_2) \right) \left((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1 \right) \Big] \\
& + \frac{G^3}{r_{12}^4} \left[\left(\frac{181m_1m_2}{16} + \frac{95m_2^2}{4} + \frac{75m_2^3}{8m_1} \right) ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_1) \right. \\
& - \left. \left(\frac{21m_1^2}{2} + \frac{473m_1m_2}{16} + \frac{63m_2^2}{4} \right) ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1) \right] + (1 \leftrightarrow 2). \tag{138}
\end{aligned}$$

This Hamiltonian is formally at 3PN but for maximally rotating objects the post-Newtonian order goes up to 3.5PN. Recently in [85, 86] the NNLO spin-orbit contributions to the acceleration and spin-precession in harmonic gauge were calculated and agreement with the equations of motion following from our Hamiltonian was found.⁵ From a combinatorial point of view there are 66 algebraically different possible contributions to the Hamiltonian for each object (written in terms of the canonical spin tensor), but 24 of them do not appear in the canonical representation used here.

7.2 Next-to-next-to-leading Order Spin(1)-Spin(2) Hamiltonian

Also the spin(1)-spin(2) Hamiltonian has an electromagnetic counterpart. It is the gravitational analogue to e.g. the coupling between electron spin and spin of the atomic nucleus, responsible for the hyperfine structure in the electromagnetic spectrum. Of course in our case the spin(1)-spin(2) interaction leads to the modulation of gravitational waves but does not lead to a hyperfine structure in the emitted atomic electromagnetic spectrum. The result for the NNLO spin(1)-spin(2) Hamiltonian reads

$$\begin{aligned}
H_{SS}^{\text{NNLO}} = & \frac{G}{r_{12}^3} \left[\frac{((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1)((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2)}{16m_1^2m_2^2} - \frac{9((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2)(\mathbf{n}_{12} \hat{\mathbf{p}}_1)}{8m_1^2m_2^2} \right. \\
& - \frac{3((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1)((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2)(\mathbf{n}_{12} \hat{\mathbf{p}}_1)}{2m_1^2m_2^2} \\
& + ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_2) \left(\frac{9\hat{\mathbf{p}}_1^2}{8m_1^4} + \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{4m_1^2m_2^2} - \frac{3\hat{\mathbf{p}}_2^2}{4m_1^2m_2^2} \right) \\
& + ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_2) \left(-\frac{3\hat{\mathbf{p}}_1^2}{2m_1^3m_2} + \frac{3(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{4m_1^2m_2^2} \right. \\
& - \left. \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1^2m_2^2} \right) + ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2) \left(\frac{3\hat{\mathbf{p}}_1^2}{16m_1^3m_2} \right. \\
& - \left. \frac{3(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{16m_1^2m_2^2} - \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{16m_1^2m_2^2} \right) \\
& + (\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_2) \left(\frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{4m_1^2m_2^2} - \frac{\hat{\mathbf{p}}_2^2}{4m_1^2m_2^2} \right) \\
& + (\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_2 \hat{\mathbf{S}}_2) \left(-\frac{\hat{\mathbf{p}}_1^2}{4m_1^3m_2} + \frac{(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{4m_1^2m_2^2} \right) \\
& + (\hat{\mathbf{p}}_2 \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_2) \left(\frac{5\hat{\mathbf{p}}_1^2}{16m_1^3m_2} - \frac{3(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{16m_1^2m_2^2} - \frac{9(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{16m_1^2m_2^2} \right) \\
& + (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_2) \left(\frac{9(\mathbf{n}_{12} \hat{\mathbf{p}}_1)\hat{\mathbf{p}}_1^2}{8m_1^4} - \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_1^2}{4m_1^3m_2} - \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_2^2}{4m_1m_2^3} \right) \\
& + (\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) \left(-\frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_2)\hat{\mathbf{p}}_1^2}{4m_1^3m_2} - \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{4m_1^2m_2^2} + \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)\hat{\mathbf{p}}_2^2}{4m_1^2m_2^2} \right)
\end{aligned}$$

⁵In [79] there is a typo in the term $-\frac{G}{r_{12}^3} \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_1)\hat{\mathbf{p}}_2^2}{16m_1^2m_2^2} ((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1)$. The coefficient has to be $-\frac{5}{16}$ instead of $-\frac{15}{16}$. The result given here is correct. Thanks to S. Marsat for pointing this out.

$$\begin{aligned}
& - \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_2) \hat{\mathbf{p}}_2^2}{4m_1 m_2^3} \Big) + (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) \Big(- \frac{3(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)^2}{8m_1^2 m_2^2} + \frac{105(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2 (\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{16m_1^2 m_2^2} \\
& - \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2 \hat{\mathbf{p}}_1^2}{8m_1^2 m_2^2} + \frac{3\hat{\mathbf{p}}_1^2 (\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{4m_1^3 m_2} + \frac{3\hat{\mathbf{p}}_1^2 \hat{\mathbf{p}}_2^2}{16m_1^2 m_2^2} + \frac{15\hat{\mathbf{p}}_1^2 (\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1^3 m_2} \Big) \\
& + (\hat{\mathbf{S}}_1 \hat{\mathbf{S}}_2) \Big(\frac{(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)^2}{16m_1^2 m_2^2} - \frac{9(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2 \hat{\mathbf{p}}_1^2}{8m_1^4} - \frac{5(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2) \hat{\mathbf{p}}_1^2}{16m_1^3 m_2} - \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2 \hat{\mathbf{p}}_1^2}{8m_1^2 m_2^2} \\
& - \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2 (\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{16m_1^2 m_2^2} + \frac{3\hat{\mathbf{p}}_1^2 \hat{\mathbf{p}}_2^2}{16m_1^2 m_2^2} + \frac{3\hat{\mathbf{p}}_1^2 (\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1^3 m_2} \\
& + \frac{9(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{16m_1^2 m_2^2} \Big) \Big] \\
& + \frac{G^2}{r_{12}^4} \Big[((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_2) \Big(\frac{12}{m_1} + \frac{9m_2}{m_1^2} \Big) \\
& - \frac{81}{4m_1} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_2) - \frac{27}{4m_1} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2) \\
& - \frac{5}{2m_1} (\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_2 \hat{\mathbf{S}}_2) + \frac{29}{8m_1} (\hat{\mathbf{p}}_2 \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_2) - \frac{21}{8m_1} (\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_2) \\
& + (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_2) \Big\{ \Big(\frac{33}{2m_1} + \frac{9m_2}{m_1^2} \Big) (\mathbf{n}_{12} \hat{\mathbf{p}}_1) - \Big(\frac{14}{m_1} + \frac{29}{2m_2} \Big) (\mathbf{n}_{12} \hat{\mathbf{p}}_2) \Big\} \\
& + (\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) \Big\{ \frac{4}{m_1} (\mathbf{n}_{12} \hat{\mathbf{p}}_1) - \Big(\frac{11}{m_1} + \frac{11}{m_2} \Big) (\mathbf{n}_{12} \hat{\mathbf{p}}_2) \Big\} \\
& + (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) \Big\{ - \frac{12}{m_1} (\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2 - \frac{10}{m_1} \hat{\mathbf{p}}_1^2 + \frac{37}{4m_1} (\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2) \\
& + \frac{255}{4m_1} (\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2) \Big\} + (\hat{\mathbf{S}}_1 \hat{\mathbf{S}}_2) \Big\{ - \Big(\frac{25}{2m_1} + \frac{9m_2}{m_1^2} \Big) (\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2 + \frac{49}{8m_1} \hat{\mathbf{p}}_1^2 \\
& + \frac{35}{4m_1} (\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2) - \frac{43}{8m_1} (\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2) \Big\} \Big] \\
& + \frac{G^3}{r_{12}^5} \Big[- (\hat{\mathbf{S}}_1 \hat{\mathbf{S}}_2) \Big(\frac{63}{4} m_1^2 + \frac{145}{8} m_1 m_2 \Big) + (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) \Big(\frac{105}{4} m_1^2 + \frac{289}{8} m_1 m_2 \Big) \Big] \\
& + (1 \leftrightarrow 2), \tag{139}
\end{aligned}$$

This Hamiltonian is formally also at 3PN but for maximally rotating objects the post-Newtonian order goes up to 4PN. Notice that from a combinatorial point of view there are 167 algebraically different possible contributions to the Hamiltonian for all objects (written in terms of the canonical spin tensor), but 75 of them do not appear in the canonical representation used here.

7.3 Hamiltonians in Center-Of-Mass Frame

For later computations of, e.g., the mentioned orbital parametrizations of a binary system it is convenient to provide the Hamiltonians in the center-of-mass frame ($\hat{\mathbf{p}}_1 = -\hat{\mathbf{p}}_2 = \hat{\mathbf{p}}$). In this frame in dimensionless quantities (see e.g. [96, 97] for rescaling) they are given by

$$\begin{aligned}
H_{\text{COM SO}}^{\text{NNLO}} &= \frac{1}{4r_{12}^5} \Big[21\sqrt{1-4\eta}(\eta+1)(\mathbf{L} \boldsymbol{\Delta}) + \frac{1}{2}(-2\eta^2 + 33\eta + 42)(\mathbf{L} \boldsymbol{\Sigma}) \Big] \\
&+ \frac{\eta}{32r_{12}^4} \Big[-\sqrt{1-4\eta}((256+45\eta)(\mathbf{n}_{12} \hat{\mathbf{p}})^2 + (314+39\eta)\hat{\mathbf{p}}^2)(\mathbf{L} \boldsymbol{\Delta}) \\
&+ ((-256+275\eta)(\mathbf{n}_{12} \hat{\mathbf{p}})^2 + (-206+73\eta)\hat{\mathbf{p}}^2)(\mathbf{L} \boldsymbol{\Sigma}) \Big] \\
&+ \frac{\eta}{32r_{12}^3} \Big[\sqrt{1-4\eta}(15(\mathbf{n}_{12} \hat{\mathbf{p}})^4 + 3(9\eta-4)(\mathbf{n}_{12} \hat{\mathbf{p}})^2 \hat{\mathbf{p}}^2
\end{aligned}$$

$$\begin{aligned}
& + 2(22\eta - 9)(\hat{\mathbf{p}}^2)^2(\mathbf{L} \Delta) - (15(2\eta - 1)(\mathbf{n}_{12} \hat{\mathbf{p}})^4 \\
& + 3(6\eta^2 - 11\eta + 4)(\mathbf{n}_{12} \hat{\mathbf{p}})^2 \hat{\mathbf{p}}^2 + 2(5\eta^2 - 3\eta + 2)(\hat{\mathbf{p}}^2)^2)(\mathbf{L} \Sigma) \Big], \tag{140a}
\end{aligned}$$

$$\begin{aligned}
H_{\text{COM SS}}^{\text{NNLO}} = & \eta \left\{ \frac{1}{4r_{12}^5} \left[(79\eta + 105)(\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) - (63 + 19\eta)(\hat{\mathbf{S}}_1 \hat{\mathbf{S}}_2) \right] \right. \\
& + \frac{1}{r_{12}^4} \left[- \left(\frac{303}{4}\eta(\mathbf{n}_{12} \hat{\mathbf{p}})^2 + \left(\frac{125}{4}\eta + 9 \right) \hat{\mathbf{p}}^2 \right) (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) \right. \\
& \quad \left(- \left(18 + \frac{25}{4}\eta \right) (\mathbf{n}_{12} \hat{\mathbf{p}})^2 + \left(9 + \frac{47}{2}\eta \right) \hat{\mathbf{p}}^2 \right) (\hat{\mathbf{S}}_1 \hat{\mathbf{S}}_2) \\
& \quad - \frac{9}{4}(7\eta + 4)(\hat{\mathbf{p}} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}} \hat{\mathbf{S}}_2) \\
& \quad + \left(34\eta + \frac{27}{2} \right) (\mathbf{n}_{12} \hat{\mathbf{p}})((\hat{\mathbf{p}} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) + (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}} \hat{\mathbf{S}}_2)) \\
& \quad \left. + \frac{3}{2}\sqrt{1 - 4\eta}(\eta + 3)(\mathbf{n}_{12} \hat{\mathbf{p}})((\hat{\mathbf{p}} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) - (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}} \hat{\mathbf{S}}_2)) \right] \\
& + \frac{1}{r_{12}^3} \left[\frac{1}{8} \left(105\eta^2(\mathbf{n}_{12} \hat{\mathbf{p}})^4 + 15\eta(3\eta - 2)(\mathbf{n}_{12} \hat{\mathbf{p}})^2 \hat{\mathbf{p}}^2 \right. \right. \\
& \quad \left. + \frac{3}{2}(10\eta^2 + 13\eta - 6)(\hat{\mathbf{p}}^2)^2 \right) (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) \\
& \quad + \frac{1}{8} \left(-3(8\eta^2 - 37\eta + 12)(\mathbf{n}_{12} \hat{\mathbf{p}})^2 \hat{\mathbf{p}}^2 + (7\eta^2 - 23\eta + 9)(\hat{\mathbf{p}}^2)^2 \right) (\hat{\mathbf{S}}_1 \hat{\mathbf{S}}_2) \\
& \quad + \frac{1}{4} \left(9\eta^2(\mathbf{n}_{12} \hat{\mathbf{p}})^2 + \frac{1}{2}(4\eta^2 + 25\eta - 9)\hat{\mathbf{p}}^2 \right) (\hat{\mathbf{p}} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}} \hat{\mathbf{S}}_2) \\
& \quad - \frac{3}{8} \left(+15\eta^2(\mathbf{n}_{12} \hat{\mathbf{p}})^2 + \frac{1}{2}(10\eta^2 + 21\eta - 9)\hat{\mathbf{p}}^2 \right) (\mathbf{n}_{12} \hat{\mathbf{p}}) \\
& \quad \quad \times ((\hat{\mathbf{p}} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) + (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}} \hat{\mathbf{S}}_2)) \\
& \quad \left. + \frac{9}{16}\sqrt{1 - 4\eta}(1 - 2\eta)(\mathbf{n}_{12} \hat{\mathbf{p}})((\hat{\mathbf{p}} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) - (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}} \hat{\mathbf{S}}_2)) \right] \Big\}. \tag{140b}
\end{aligned}$$

There $\Delta = \hat{\mathbf{S}}_1 - \hat{\mathbf{S}}_2$ and $\Sigma = \hat{\mathbf{S}}_1 + \hat{\mathbf{S}}_2$ the differences and sums of the spin vectors and $\mathbf{L}$ is the orbital angular momentum $\mathbf{L} = r_{12}\mathbf{n}_{12} \times \hat{\mathbf{p}}$.

8 Kinematical Consistency: The Approximate Poincaré Algebra

For a space-time which is asymptotically flat the Poincaré algebra must be fulfilled at spacial infinity, e.g. [189]. The generators of the Poincaré algebra can be expressed in terms of the canonical variables describing the physical system, i.e. matter variables like linear momenta, position variables or spins. Also propagating field degrees of freedom enter the generators of the Poincaré algebra. Throughout this section we set $d = 3$. The relations between the generators are given by

$$\{P_i, H\} = 0, \quad \{J_i, H\} = 0, \tag{141a}$$

$$\{J_i, P_j\} = \epsilon_{ijk}P_k, \quad \{J_i, J_j\} = \epsilon_{ijk}J_k, \tag{141b}$$

$$\{J_i, G_j\} = \epsilon_{ijk}G_k, \tag{141c}$$

$$\{G_i, H\} = P_i, \tag{141d}$$

$$\{G_i, P_j\} = c^{-2}\delta_{ij}H, \tag{141e}$$

$$\{G_i, G_j\} = -c^{-2}\epsilon_{ijk}J_k, \tag{141f}$$

where $\mathbf{P}$ is the total linear momentum, J^{ij} is the total angular momentum tensor and $J_i = \frac{1}{2}\epsilon_{ijk}J^{jk}$ the associated dual vector, $\mathbf{G}$ is the center-of-mass vector and H the Hamiltonian of the physical system. Total linear momentum $\mathbf{P}$ and total angular momentum $J^{ij} = -J^{ji}$ are given by

$$\mathbf{P} = \sum_a \hat{\mathbf{p}}_a, \quad J^{ij} = \sum_a \left[\hat{z}_a^i \hat{p}_a^j - \hat{z}_a^j \hat{p}_a^i + \hat{S}_{a(i)(j)} \right], \quad (142)$$

see also, e.g., [30, 51]. For the contributions of the propagating field degrees of freedom see, e.g., [76, 77]. However, these contributions can be dropped within $\mathbf{P}$ and J^{ij} here as we are considering the *conservative* matter-only Hamiltonian instead of the ADM Hamiltonian (the latter still depends on the canonical field variables).

8.1 General Considerations for a Center-Of-Mass Vector Ansatz

As in [30, 51, 79] we use an ansatz for the center-of-mass vectors $\mathbf{G}$ at next-to-next-to-leading order (at lower orders it is also possible to directly calculate $\mathbf{G}$ from certain integrals). For constructing the center-of-mass vectors one has to consider the irreducible algebraic quantities which can be generated from $\hat{S}_{a(i)(j)}$, $\hat{p}_{a i}$ and n_{ab}^i . Since the Newtonian center-of-mass vector

$$\mathbf{G}_N = \sum_a m_a \hat{\mathbf{z}}_a, \quad (143)$$

is at c^{-2} and the Newtonian Hamiltonian

$$H_N = \sum_a \frac{\hat{\mathbf{p}}_a^2}{2m_a} - \sum_a \sum_{b \neq a} \frac{Gm_a m_b}{2r_{ab}}, \quad (144)$$

is at c^{-4} the higher order corrections to the center-of-mass vector are also one post-Newtonian order below the appropriate Hamiltonian. Thus the momentum and G powers there are also reduced.

Let us demonstrate these considerations at point-mass level: The Newtonian Hamiltonian has only p^2 terms at G^0 [which could be $\hat{\mathbf{p}}_1^2$, $\hat{\mathbf{p}}_2^2$, $(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2$, $(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2$, and $(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)^6$ and p^0 terms at G^1 (which is only one term). At 1PN there appear p^4 terms at G^0 , p^2 terms at G^1 and p^0 at G^2 . At 3PN there are p^8 terms at G^0 and p^0 terms at G^4 . The center-of-mass vectors belonging to the Hamiltonians above have the following momentum powers emerging there: The Newtonian center-of-mass vector mentioned above has p^0 at G^0 , the 1PN one contains p^2 at G^0 and p^0 at G^1 and at 3PN level it contains p^6 at G^0 and p^0 at G^3 .

Now we will discuss how to construct the linear-in-spin corrections for the center-of-mass vectors. Symbolically they can be written in the form

$$\mathbf{G}_{\text{SO}} = \text{SO-scalar} \cdot \text{PM-vector} + \text{PM-scalar} \cdot \text{SO-vector}, \quad (145)$$

$$\mathbf{G}_{\text{SS}} = S_1 S_2\text{-scalar} \cdot \text{PM-vector} + \text{SO-scalar} \cdot \text{SO-vector} + \text{PM-scalar} \cdot S_1 S_2\text{-vector}. \quad (146)$$

Notice that we are formally working in generic dimension, where a *spin-vector* can not be defined. As the Poincare-Algebra must also hold in generic dimensions, it must be possible to construct the center-of-mass vector in terms of the *spin-tensor*. This is fortunate, as identities such as (6.1) in [85] would complicate the situation if one is forced to work with a spin vector in $d = 3$.

Let us now summarize the vector quantities which can be built at certain spin-levels and may be used to construct the center-of-mass vectors.

The mentioned vectors must perhaps be multiplied by scalar quantities, see (145) and (146). If the number of momentum variables in the spin-orbit or spin(1)-spin(2) scalars given in the following Table 2 is not sufficient for the appropriate G -order they have to be filled up by the point-mass scalars, namely the linear momentum powers.

Also important is that every spin is counted like a linear momentum because they have the same c^{-1} -order, see 3.1. This means the formal 3PN spin-orbit and spin(1)-spin(2) Hamiltonians have only contributions up to G^3 (G^4 contributions cannot contain any spins since they are momentum-independent for

⁶Although only $\hat{\mathbf{p}}_1^2$ and $\hat{\mathbf{p}}_2^2$ are appearing at the Newtonian order. This discussion should only provide some idea of appearing momentum powers at certain post-Newtonian orders.

Vector	Irreducible Quantities
point-mass (PM)	$\hat{\mathbf{z}}_a, \hat{\mathbf{p}}_a, \mathbf{n}_{ab}$
spin-orbit (SO) (for $\hat{S}_{a(i)(j)}$)	$n_{ab}^j \hat{S}_{a(i)(j)} \hat{p}_{a j} \hat{S}_{a(i)(j)}, \hat{p}_{b j} \hat{S}_{a(i)(j)}$
spin(a)-spin(b) ($S_a S_b$)	$n_{ab}^k \hat{S}_{a(k)(j)} \hat{S}_{b(i)(j)}, \hat{p}_{a k} \hat{S}_{a(k)(j)} \hat{S}_{b(i)(j)}, \hat{p}_{b k} \hat{S}_{a(k)(j)} \hat{S}_{b(i)(j)}$

Table 1: Vector quantities at certain spin levels

Scalar	Irreducible Quantities
PM	linear momentum powers p^n
SO	$(n_{ab}^i \hat{p}_{a j} \hat{S}_{a(i)(j)}), (n_{ab}^i \hat{p}_{b j} \hat{S}_{a(i)(j)}), (\hat{p}_{a i} \hat{p}_{b j} \hat{S}_{a(i)(j)})$
$S_1 S_2$	$(\hat{p}_{1 i} \hat{p}_{2 j} \hat{S}_{1(i)(j)})(\hat{p}_{1 i} \hat{p}_{2 j} \hat{S}_{2(i)(j)}), (n_{12}^i \hat{p}_{1 j} \hat{S}_{1(i)(j)})(n_{12}^i \hat{p}_{1 j} \hat{S}_{2(i)(j)}),$ $(n_{12}^i \hat{p}_{1 j} \hat{S}_{1(i)(j)})(n_{12}^i \hat{p}_{2 j} \hat{S}_{2(i)(j)}), (\hat{p}_{1 i} \hat{p}_{2 j} \hat{S}_{1(i)(j)})(\hat{p}_{1 i} \hat{p}_{2 j} \hat{S}_{2(i)(j)}),$ $(n_{12}^i \hat{p}_{1 j} \hat{S}_{1(i)(j)})(n_{12}^i \hat{p}_{1 j} \hat{S}_{2(i)(j)}), (n_{12}^i \hat{p}_{1 j} \hat{S}_{1(i)(j)})(n_{12}^i \hat{p}_{2 j} \hat{S}_{2(i)(j)}),$ $(n_{12}^i \hat{p}_{2 j} \hat{S}_{1(i)(j)})(n_{12}^i \hat{p}_{1 j} \hat{S}_{2(i)(j)}), (n_{12}^i \hat{p}_{2 j} \hat{S}_{1(i)(j)})(n_{12}^i \hat{p}_{2 j} \hat{S}_{2(i)(j)}),$ $(\hat{S}_{1(i)(j)} \hat{S}_{2(i)(j)}), (n_{12}^i n_{12}^j \hat{S}_{1(i)(k)} \hat{S}_{2(j)(k)}),$ $(n_{12}^i \hat{p}_{1 j} \hat{S}_{1(i)(k)} \hat{S}_{2(j)(k)}), (n_{12}^i \hat{p}_{2 j} \hat{S}_{1(i)(k)} \hat{S}_{2(j)(k)}),$ $(\hat{p}_{1 i} n_{12}^j \hat{S}_{1(i)(k)} \hat{S}_{2(j)(k)}), (\hat{p}_{2 i} n_{12}^j \hat{S}_{1(i)(k)} \hat{S}_{2(j)(k)}),$ $(\hat{p}_{1 i} \hat{p}_{2 j} \hat{S}_{1(i)(k)} \hat{S}_{2(j)(k)}), (\hat{p}_{2 i} \hat{p}_{1 j} \hat{S}_{1(i)(k)} \hat{S}_{2(j)(k)}),$ $(\hat{p}_{1 i} \hat{p}_{2 j} \hat{S}_{1(i)(j)})(n_{12}^i \hat{p}_{1 j} \hat{S}_{2(i)(j)}), (\hat{p}_{1 i} \hat{p}_{2 j} \hat{S}_{1(i)(j)})(n_{12}^i \hat{p}_{2 j} \hat{S}_{2(i)(j)}),$ $(n_{12}^i \hat{p}_{1 j} \hat{S}_{1(i)(j)})(\hat{p}_{1 i} \hat{p}_{2 j} \hat{S}_{2(i)(j)}), (n_{12}^i \hat{p}_{2 j} \hat{S}_{1(i)(j)})(\hat{p}_{1 i} \hat{p}_{2 j} \hat{S}_{2(i)(j)})$

Table 2: Scalar quantities at certain spin levels

point-masses). Notice that the Hamiltonians can only be constructed from the irreducible scalar quantities given above. This is demanded by the Poincaré algebra, namely (141a) (H should be invariant under translations and rotations and thus is a scalar). The $\hat{\mathbf{z}}_a$ contribution in the center-of-mass vector can be fixed by (141e) using the lower order Hamiltonian. This Hamiltonian can always be written in the form $H = \sum_a h_a$, where the h_a are translation invariant, $\{h_a, \mathbf{P}\} = 0$. (In the post-Newtonian approximation of general relativity all Hamiltonians have such a structure that the h_a are translational invariant.) If we make an ansatz for the center-of-mass vector of the form

$$\mathbf{G} = \sum_a h_a \hat{\mathbf{z}}_a + \mathbf{Y}, \quad (147)$$

we see that ($c^{-1} = 1$)

$$\left\{ \sum_a h_a \hat{z}_a^i + Y^i, \sum_b \hat{p}_{b j} \right\} = \sum_a \left[\underbrace{\{h_a, P^j\}}_{=0} \hat{z}_a^i + h_a \delta_{ab} \delta_{ij} \right] + \underbrace{\{Y^i, P^j\}}_{=0} = \sum_a h_a \delta_{ij} = H \delta_{ij}. \quad (148)$$

Equation (148) demands that $\{Y^i, P^j\} = 0$ and so $\mathbf{Y}$ must be translational invariant. We have shown that the part of the center-of-mass vector which is not translation invariant, i.e., $\sum_a h_a \hat{\mathbf{z}}_a$, can be read off from the Hamiltonian.

From these consideration it follows that in the spin-orbit case the center-of-mass vector consists of 52 algebraic independent quantities for one object and in the spin(1)-spin(2) case there are 86 algebraic independent quantities for both objects. Notice that up to the formal 3PN level and linear order in spin all center-of-mass vectors can be fixed uniquely by using the Poincaré algebra.

8.2 Next-to-next-to-leading Order linear-in-spin Center-Of-Mass Vectors

Now we take the ansatz for the center-of-mass vector (147) where $\mathbf{Y}$ has to be constructed from the irreducible quantities given in Tables 1 and 2 with the decompositions (145) and (146), but without $\hat{\mathbf{z}}_a$ vectors

and put them into (141d) with the Hamiltonians (138) and (139) for the spin-orbit and spin(1)-spin(2) case. From this all unknown coefficients mentioned above could be fixed uniquely. The center-of-mass vector contributions given here implement the change in the binding energy of the system due to the NNLO linear-in-spin interaction Hamiltonians. This results also in context of the energy-mass equivalence in a modified gravitating mass and thus in a correction to the Newtonian center-of-mass vector $\mathbf{G}_N = \sum_a m_a \hat{\mathbf{z}}_a$, which does not take any interactions into account. The correction to the center-of-mass vector from NNLO spin-orbit interactions finally results as

$$\begin{aligned}
\mathbf{G}_{\text{SO}}^{\text{NNLO}} = & \frac{(\hat{\mathbf{p}}_1^2)^2}{16m_1^5} (\hat{\mathbf{p}}_1 \times \hat{\mathbf{S}}_1) \\
& + (\hat{\mathbf{p}}_2 \times \hat{\mathbf{S}}_1) \left[\frac{G}{r_{12}} \left(-\frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{8m_1 m_2} - \frac{(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{8m_1 m_2} \right) + \frac{G^2}{r_{12}^2} \left(-\frac{47m_1}{16} - \frac{21m_2}{8} \right) \right] \\
& + (\hat{\mathbf{p}}_1 \times \hat{\mathbf{S}}_1) \left[\frac{G}{r_{12}} \left(\frac{9m_2 \hat{\mathbf{p}}_1^2}{16m_1^3} - \frac{5\hat{\mathbf{p}}_2^2}{8m_1 m_2} \right) + \frac{G^2}{r_{12}^2} \left(\frac{57m_2}{16} + \frac{15m_2^2}{8m_1} \right) \right] \\
& + (\mathbf{n}_{12} \times \hat{\mathbf{S}}_1) \left[\frac{G}{r_{12}} \left(\frac{9(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{16m_1 m_2} + \frac{(\mathbf{n}_{12} \hat{\mathbf{p}}_2)(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{8m_1 m_2} + \frac{(\mathbf{n}_{12} \hat{\mathbf{p}}_1)\hat{\mathbf{p}}_2^2}{16m_1 m_2} \right) \right. \\
& \quad \left. + \frac{G^2}{r_{12}^2} \left(-\frac{5m_2}{8} (\mathbf{n}_{12} \hat{\mathbf{p}}_1) + \left\{ \frac{13m_1}{8} + \frac{11m_2}{4} \right\} (\mathbf{n}_{12} \hat{\mathbf{p}}_2) \right) \right] \\
& - \frac{G}{r_{12}} \hat{\mathbf{p}}_1 \frac{(\mathbf{n}_{12} \hat{\mathbf{p}}_2)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1)}{2m_1 m_2} \\
& + \frac{G}{r_{12}} \hat{\mathbf{p}}_2 \left(-\frac{(\mathbf{n}_{12} \hat{\mathbf{p}}_2)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1)\hat{\mathbf{S}}_1)}{8m_1 m_2} + \frac{(\mathbf{n}_{12} \hat{\mathbf{p}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1)}{2m_1 m_2} \right. \\
& \quad \left. - \frac{((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1)}{8m_1 m_2} \right) \\
& + \mathbf{n}_{12} \left[\frac{G}{r_{12}} \left(\left\{ \frac{m_2 \hat{\mathbf{p}}_1^2}{16m_1^3} + \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{16m_1 m_2} - \frac{3\hat{\mathbf{p}}_2^2}{16m_1 m_2} \right\} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1)\hat{\mathbf{S}}_1) \right. \right. \\
& \quad + \left\{ -\frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{2m_1 m_2} - \frac{(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{2m_1 m_2} \right\} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1) \\
& \quad \left. + \frac{13(\mathbf{n}_{12} \hat{\mathbf{p}}_2)((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1)}{8m_1 m_2} \right) \\
& \quad \left. + \frac{G^2}{r_{12}^2} \left(\left\{ \frac{m_2}{2} + \frac{5m_2^2}{4m_1} \right\} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1)\hat{\mathbf{S}}_1) + \{-2m_1 - 5m_2\} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1) \right) \right] \\
& + \frac{\hat{\mathbf{z}}_1}{r_{12}} \left[\frac{G}{r_{12}} \left(\left\{ \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{m_1 m_2} + \frac{(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{m_1 m_2} \right\} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1) \right. \right. \\
& \quad + \left\{ \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)}{4m_1^2} - \frac{2(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{m_1 m_2} \right\} ((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_1) \\
& \quad + \left\{ -\frac{5m_2 \hat{\mathbf{p}}_1^2}{8m_1^3} - \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1^2} - \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2}{2m_1 m_2} \right. \\
& \quad \left. \left. - \frac{3(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{4m_1^2} + \frac{3\hat{\mathbf{p}}_2^2}{4m_1 m_2} \right\} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1)\hat{\mathbf{S}}_1) \right) \\
& \quad \left. + \frac{G^2}{r_{12}^2} \left(\left\{ -\frac{11m_2}{2} - \frac{5m_2^2}{m_1} \right\} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1)\hat{\mathbf{S}}_1) + \left\{ 6m_1 + \frac{15m_2}{2} \right\} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2)\hat{\mathbf{S}}_2) \right) \right] \\
& + (1 \leftrightarrow 2), \tag{149}
\end{aligned}$$

and the NNLO spin(1)-spin(2) part reads

$$\mathbf{G}_{\text{SS}}^{\text{NNLO}} = \frac{G^2}{r_{12}^3} ((\mathbf{n}_{12} \times \hat{\mathbf{S}}_2) \times \hat{\mathbf{S}}_1) \left(\frac{17}{8} m_1 + m_2 \right)$$

$$\begin{aligned}
& + \frac{G}{r_{12}^2} \left[\hat{\mathbf{p}}_1 \left(-\frac{(\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_2 \hat{\mathbf{S}}_2)}{4m_1 m_2} + \frac{3(\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1 m_2} \right) \right. \\
& + (\mathbf{n}_{12} \times \hat{\mathbf{S}}_1) \left(-\frac{((\hat{\mathbf{p}}_1 \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2)}{4m_1 m_2} + \frac{3((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_2)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1 m_2} \right) \\
& - (\hat{\mathbf{p}}_1 \times \hat{\mathbf{S}}_1) \frac{((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2)}{8m_1 m_2} - (\hat{\mathbf{p}}_2 \times \hat{\mathbf{S}}_1) \frac{((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_2)}{4m_1 m_2} \\
& - ((\hat{\mathbf{p}}_1 \times \hat{\mathbf{S}}_2) \times \hat{\mathbf{S}}_1) \frac{(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1 m_2} - ((\hat{\mathbf{p}}_2 \times \hat{\mathbf{S}}_2) \times \hat{\mathbf{S}}_1) \frac{(\mathbf{n}_{12} \hat{\mathbf{p}}_1)}{4m_1 m_2} \Big] \\
& + \frac{\hat{\mathbf{z}}_1}{r_{12}} \left(\frac{2G^2(2m_1 + m_2)}{r_{12}^3} \left[(\hat{\mathbf{S}}_1 \hat{\mathbf{S}}_2) - 2(\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) \right] \right. \\
& + \frac{G}{r_{12}^2} \left[-\frac{3((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_2)}{2m_1^2} \right. \\
& + \frac{3((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_2)}{2m_1 m_2} + \frac{3((\mathbf{n}_{12} \times \hat{\mathbf{p}}_1) \hat{\mathbf{S}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2)}{8m_1 m_2} \\
& - \frac{(\hat{\mathbf{p}}_2 \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_2)}{8m_1 m_2} + \frac{(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_2 \hat{\mathbf{S}}_2)}{4m_1 m_2} + \frac{3(\hat{\mathbf{p}}_2 \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2)(\mathbf{n}_{12} \hat{\mathbf{p}}_1)}{2m_1 m_2} \\
& - \frac{3(\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_2)(\mathbf{n}_{12} \hat{\mathbf{p}}_1)}{2m_1^2} + \frac{3(\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\hat{\mathbf{p}}_2 \hat{\mathbf{S}}_2)(\mathbf{n}_{12} \hat{\mathbf{p}}_1)}{4m_1 m_2} \\
& + \frac{3(\hat{\mathbf{p}}_1 \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1 m_2} \\
& - (\mathbf{n}_{12} \hat{\mathbf{S}}_1)(\mathbf{n}_{12} \hat{\mathbf{S}}_2) \left\{ \frac{15(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1 m_2} + \frac{3(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{4m_1 m_2} \right\} \\
& \left. \left. + (\hat{\mathbf{S}}_1 \hat{\mathbf{S}}_2) \left\{ \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)^2}{2m_1^2} - \frac{3(\mathbf{n}_{12} \hat{\mathbf{p}}_1)(\mathbf{n}_{12} \hat{\mathbf{p}}_2)}{4m_1 m_2} + \frac{(\hat{\mathbf{p}}_1 \hat{\mathbf{p}}_2)}{8m_1 m_2} \right\} \right] \right) + (1 \leftrightarrow 2). \tag{150}
\end{aligned}$$

From this the boost vector $\mathbf{K} = \mathbf{G} - t\mathbf{P}$ can be obtained, which explicitly depends on time t .

Notice that in (149) (in contrast to (150)) there appears a one-particle term without G factor. It comes from the displacement of the center-of-mass due to the rotation and the resulting special relativistic Lorentz contractions of different parts of the object (which has to have a finite size). Further discussions about this issue can be found in [76] which are clarified graphically in particular in Fig. 1 therein. In (150) there is no term without G factor because all interactions between two spins are transmitted by the gravitational field in general relativity.

9 Test-Spin near Kerr Black Hole

In the last section we checked whether our results are compatible with kinematical restrictions of the Poincaré algebra. Here we derive an exact test-spin Hamiltonian to compare our results (138) and (139) with. In the following subsections we restrict ourselves to $d = 3$, because in the test-spin case there are only delta integrals to be evaluated. A partial check of our Hamiltonians against the test-spin case is contained in [74] for the case of aligned spins.

There are various approaches to calculate the motion of a test-spin near a Kerr black hole (see, e.g., [190] and references therein). Since in the counting used in [190] the NNLO spin(1)-spin(2) interaction is at 4PN and therefore not considered therein, one needs to calculate the spin(1)-spin(2) contributions from

$$H_{\text{Testspin}} = -\hat{A}^{k\ell} e_{(m)k} e_{\ell,0}^{(m)} + \int d^3x [\mathcal{H}^{\text{matter}} N - \mathcal{H}_i^{\text{matter}} N^i], \tag{151}$$

where N , N_i , the framefield $e_{(m)k}$, and the implicit appearing metric provide the exterior gravitational field, and $\mathcal{H}^{\text{matter}}$ and $\mathcal{H}_i^{\text{matter}}$ represent the test-spin moving in it [76]. Note that the framefield in the first term has to be evaluated at the position of the test-spin.

Since all spin dependencies of the metric are at least quadratic in the Kerr spin, and the only contribution which is linear in Kerr spin comes from the shift vector, the three-dimensional part of the metric and the lapse are identical with the appropriate components of the isotropic Schwarzschild metric which also comes from the spinless limit of the Kerr metric. The shift is given by the expressions in [60, Eq. (54)].

The metric components generated by particle ‘1’ are given by

$$g_{00} = - \left(\frac{1 - \frac{m_1}{2r_1}}{1 + \frac{m_1}{2r_1}} \right)^2, \quad (152)$$

$$g_{ij} = \left(1 + \frac{m_1}{2r_1} \right)^4 \delta_{ij}, \quad (153)$$

$$g_{0i} = \frac{2m_1 n_1^k \hat{a}_{1(i)(k)}}{r_1^2 \left(1 + \frac{m_1}{2r_1} \right)^2}. \quad (154)$$

It is well-known that the metric components can be rewritten into a three-dimensional metric on the spatial hypersurface, lapse N , and shift $N^i = \gamma^{ij} N_j$, using (2) and (3). So one can immediately see that

$$\gamma_{ij} = g_{ij} = \left(1 + \frac{m_1}{2r_1} \right)^4 \delta_{ij}, \quad (155)$$

$$\gamma^{ij} = \left(1 + \frac{m_1}{2r_1} \right)^{-4} \delta_{ij}, \quad (156)$$

$$N = \frac{1 - \frac{m_1}{2r_1}}{1 + \frac{m_1}{2r_1}}, \quad (157)$$

with the squares of the shift neglected, because they are quadratic in the Kerr spin,

$$N_i = \frac{2m_1 n_1^k \hat{a}_{1(i)(k)}}{r_1^2 \left(1 + \frac{m_1}{2r_1} \right)^2} = -2m_1 \hat{a}_{1(i)(k)} \left(\frac{1}{r_1 \left(1 + \frac{m_1}{2r_1} \right)} \right)_{,k}, \quad (158)$$

see [60]. Here $\hat{a}_{1(i)(j)} = \hat{S}_{1(i)(j)}/m_1$ is the Kerr spin belonging to the black hole located at position ‘1’. Note that to linear order in spin

$$\pi^{ij} = \frac{6m_1 n_1^k n_1^{(i} \hat{a}_{1(j))(k)}}{r_1^3 \left(1 + \frac{m_1}{2r_1} \right)^4}, \quad (159)$$

(calculated from the inverse metric and the three-dimensional Christoffel symbols, see [60, Eq. (65)]) fulfills the ADM gauge condition, so no further coordinate shift from quasi isotropic coordinates to another coordinate system is necessary. Due to the symmetric framefield gauge $e_{(i)j} = \sqrt{\gamma_{ij}}$, the framefield is given by (101)

$$e_{(i)j} = \left(1 + \frac{m_1}{2r_1} \right)^2 \delta_{ij}. \quad (160)$$

For the test-spin in a Kerr field it is sufficient to calculate (151) where the sources are given by (94) and (95). Since the metric and the framefield are proportional to δ_{ij} many terms vanish in the source. The only terms remaining are

$$\mathcal{H}^{\text{matter}} = \sum_a \left[-n \hat{p}_a \delta_a - \frac{1}{2} \frac{\hat{S}_{a li} \hat{p}_{a j}}{n \hat{p}_a} \gamma^{kl} \gamma^{ij}{}_{,k} \delta_a - \left(\frac{\hat{p}_{a l}}{m_a - n \hat{p}_a} \gamma^{ij} \gamma^{kl} \hat{S}_{a jk} \delta_a \right)_{,i} \right], \quad (161)$$

$$\mathcal{H}_i^{\text{matter}} = \sum_a \left[\hat{p}_{a i} \delta_a + \frac{1}{2} \left(\gamma^{jk} \hat{S}_{a ik} \delta_a + \gamma^{jk} \gamma^{\ell p} \frac{2 \hat{p}_a \ell \hat{p}_{a (i} \hat{S}_{a k) p}}{n \hat{p}_a (m_a - n \hat{p}_a)} \delta_a \right)_{,j} \right]. \quad (162)$$

Inserting sources, metric components and framefield, and evaluating them at the testspin location gives the exact result

$$\begin{aligned}
H_{\text{Testspin}}^{\text{Kerr}} &\stackrel{\mathcal{O}(a^1)}{=} \frac{1 - \frac{m_1}{2r_{12}}}{1 + \frac{m_1}{2r_{12}}} \sqrt{m_2^2 + \frac{\hat{\mathbf{p}}_2^2}{\left(1 + \frac{m_1}{2r_{12}}\right)^4}} \\
&- \frac{m_1((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{r_{12}^2 \left(1 + \frac{m_1}{2r_{12}}\right)^6} \left[\frac{1 - \frac{m_1}{2r_{12}}}{\sqrt{m_2^2 + \frac{\hat{\mathbf{p}}_2^2}{\left(1 + \frac{m_1}{2r_{12}}\right)^4}}} + \frac{1}{m_2 + \sqrt{m_2^2 + \frac{\hat{\mathbf{p}}_2^2}{\left(1 + \frac{m_1}{2r_{12}}\right)^4}}} \right] \\
&+ \frac{2m_1((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{a}}_1)}{r_{12}^2 \left(1 + \frac{m_1}{2r_{12}}\right)^6} + \frac{m_1}{r_{12}^3 \left(1 + \frac{m_1}{2r_{12}}\right)^7} \left[- \left(1 - \frac{5m_1}{2r_{12}}\right) (\hat{\mathbf{a}}_1 \cdot \hat{\mathbf{S}}_2) \right. \\
&- 3 \left(1 - \frac{m_1}{2r_{12}}\right) \left\{ -(\mathbf{n}_{12} \cdot \hat{\mathbf{a}}_1)(\mathbf{n}_{12} \cdot \hat{\mathbf{S}}_2) \right. \\
&\left. \left. + \frac{((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{a}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2) - (\mathbf{n}_{12} \cdot \hat{\mathbf{p}}_2)^2 (\hat{\mathbf{a}}_1 \cdot \hat{\mathbf{S}}_2) + (\mathbf{n}_{12} \cdot \hat{\mathbf{p}}_2)(\hat{\mathbf{p}}_2 \cdot \hat{\mathbf{a}}_1)(\mathbf{n}_{12} \cdot \hat{\mathbf{S}}_2)}{\left(m_2 + \sqrt{m_2^2 + \frac{\hat{\mathbf{p}}_2^2}{\left(1 + \frac{m_1}{2r_{12}}\right)^4}}\right) \sqrt{m_2^2 + \frac{\hat{\mathbf{p}}_2^2}{\left(1 + \frac{m_1}{2r_{12}}\right)^4}} \left(1 + \frac{m_1}{2r_{12}}\right)^4} \right\} \right], \tag{163}
\end{aligned}$$

which leads after a post-Newtonian expansion (the post-Newtonian order given in the subscript is a formal one)

$$\begin{aligned}
H_{\text{Testspin}, \leq 3\text{PN}}^{\text{Kerr}} &\stackrel{\mathcal{O}(a^1)}{\approx} m_2 c^2 + \left(\frac{\hat{\mathbf{p}}_2^2}{2m_2} - \frac{m_1 m_2}{r_{12}} \right) + c^{-2} \left(-\frac{(\hat{\mathbf{p}}_2^2)^2}{8m_2^3} - \frac{3m_1 \hat{\mathbf{p}}_2^2}{2m_2 r_{12}} \right. \\
&+ \frac{m_1}{r_{12}^2} \left[\frac{m_1 m_2}{2} - 2((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{a}}_1) - \frac{3((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{2m_2} \right] \\
&+ \frac{m_1}{r_{12}^3} \left[-(\hat{\mathbf{a}}_1 \cdot \hat{\mathbf{S}}_2) + 3(\mathbf{n}_{12} \cdot \hat{\mathbf{a}}_1)(\mathbf{n}_{12} \cdot \hat{\mathbf{S}}_2) \right] \Big) \\
&+ c^{-4} \left(\frac{(\hat{\mathbf{p}}_2^2)^3}{16m_2^5} + \frac{5m_1(\hat{\mathbf{p}}_2^2)^2}{8m_2^3 r_{12}} + \frac{m_1 \hat{\mathbf{p}}_2^2}{m_2 r_{12}^2} \left[\frac{5m_1}{2} + \frac{5((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{8m_2^2} \right] \right. \\
&+ \frac{m_1}{r_{12}^3} \left[-\frac{m_1^2 m_2}{4} + m_1 \left(6((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{a}}_1) + \frac{5((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{m_2} \right) \right. \\
&- \frac{3((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{a}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{2m_2^2} + \frac{3(\mathbf{n}_{12} \cdot \hat{\mathbf{p}}_2)^2 (\hat{\mathbf{a}}_1 \cdot \hat{\mathbf{S}}_2)}{2m_2^2} \\
&- \frac{3(\mathbf{n}_{12} \cdot \hat{\mathbf{p}}_2)(\hat{\mathbf{p}}_2 \cdot \hat{\mathbf{a}}_1)(\mathbf{n}_{12} \cdot \hat{\mathbf{S}}_2)}{2m_2^2} \Big] + \frac{6m_1^2}{r_{12}^4} \left[(\hat{\mathbf{a}}_1 \cdot \hat{\mathbf{S}}_2) - 2(\mathbf{n}_{12} \cdot \hat{\mathbf{a}}_1)(\mathbf{n}_{12} \cdot \hat{\mathbf{S}}_2) \right] \Big) \\
&+ c^{-6} \left(-\frac{5(\hat{\mathbf{p}}_2^2)^4}{128m_2^7} - \frac{7m_1(\hat{\mathbf{p}}_2^2)^3}{16m_2^5 r_{12}} - \frac{m_1(\hat{\mathbf{p}}_2^2)^2}{m_2^3 r_{12}^2} \left[\frac{27m_1}{16} + \frac{7((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{16m_2^2} \right] \right. \\
&+ \frac{m_1 \hat{\mathbf{p}}_2^2}{m_2 r_{12}^3} \left[-\frac{25m_1^2}{8} - \frac{27m_1((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{8m_2^2} + \frac{9((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{a}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{8m_2^3} \right. \\
&+ \frac{9(\mathbf{n}_{12} \cdot \hat{\mathbf{p}}_2)(\hat{\mathbf{p}}_2 \cdot \hat{\mathbf{a}}_1)(\mathbf{n}_{12} \cdot \hat{\mathbf{S}}_2)}{8m_2^3} - \frac{9(\mathbf{n}_{12} \cdot \hat{\mathbf{p}}_2)^2 (\hat{\mathbf{a}}_1 \cdot \hat{\mathbf{S}}_2)}{8m_2^3} \Big] + \frac{m_1^2}{r_{12}^4} \left[\frac{m_1^2 m_2}{8} \right. \\
&+ m_1 \left(-\frac{21((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{a}}_1)}{2} - \frac{75((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{8m_2} \right) \\
&\left. \left. + \frac{9((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{a}}_1)((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \cdot \hat{\mathbf{S}}_2)}{m_2^2} + \frac{9(\mathbf{n}_{12} \cdot \hat{\mathbf{p}}_2)(\hat{\mathbf{p}}_2 \cdot \hat{\mathbf{a}}_1)(\mathbf{n}_{12} \cdot \hat{\mathbf{S}}_2)}{m_2^2} \right) \right]
\end{aligned}$$

$$- \frac{9(\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2 (\hat{\mathbf{a}}_1 \hat{\mathbf{S}}_2)}{m_2^2} \Big] + \frac{m_1^3}{r_{12}^5} \left[-\frac{63}{4} (\hat{\mathbf{a}}_1 \hat{\mathbf{S}}_2) + \frac{105}{4} (\mathbf{n}_{12} \hat{\mathbf{a}}_1) (\mathbf{n}_{12} \hat{\mathbf{S}}_2) \right] \Big), \quad (164)$$

in full agreement with the test-spin limit of the full point-mass Hamiltonian, the spin-orbit Hamiltonian and the spin(1)-spin(2) Hamiltonian up to and including the formal 3PN order. For later checks we provide also the expressions at formal 4PN order. The test-spin limit for the next-to-next-to-next-to-leading order (NNNLO) spin-orbit interaction is given by

$$H_{\text{Testspin,4PN,SO}}^{\text{Kerr}} = \left(\frac{45}{128} \frac{m_1 (\hat{\mathbf{p}}_2^2)^3}{m_2^7 r_{12}^2} + \frac{13}{4} \frac{m_1^2 (\hat{\mathbf{p}}_2^2)^2}{m_2^5 r_{12}^3} + \frac{315}{32} \frac{m_1^3 \hat{\mathbf{p}}_2^2}{m_2^3 r_{12}^4} + \frac{105}{8} \frac{m_1^4}{m_2 r_{12}^5} \right) ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2) + 14 \frac{m_1^4}{r_{12}^5} ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{a}}_1), \quad (165)$$

and for the NNNLO spin(1)-spin(2) interaction by

$$H_{\text{Testspin,4PN,SS}}^{\text{Kerr}} = \left(-\frac{15}{16} \frac{m_1 (\hat{\mathbf{p}}_2^2)^2}{m_2^6 r_{12}^3} - 9 \frac{m_1^2 \hat{\mathbf{p}}_2^2}{m_2^4 r_{12}^4} - \frac{231}{8} \frac{m_1^3}{m_2^2 r_{12}^5} \right) [((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{a}}_1) ((\mathbf{n}_{12} \times \hat{\mathbf{p}}_2) \hat{\mathbf{S}}_2) + (\mathbf{n}_{12} \hat{\mathbf{p}}_2) (\hat{\mathbf{p}}_2 \hat{\mathbf{a}}_1) (\mathbf{n}_{12} \hat{\mathbf{S}}_2) - (\mathbf{n}_{12} \hat{\mathbf{p}}_2)^2 (\hat{\mathbf{a}}_1 \hat{\mathbf{S}}_2)] + \frac{14m_1^4}{r_{12}^6} \left(2(\hat{\mathbf{a}}_1 \hat{\mathbf{S}}_2) - 3(\mathbf{n}_{12} \hat{\mathbf{a}}_1) (\mathbf{n}_{12} \hat{\mathbf{S}}_2) \right). \quad (166)$$

Notice that the formal 3PN spin(1)-spin(2) test-spin contributions were not given in [190] (in their counting rules they would be at 4PN level). Further notice that there are no contributions coming from the first term in (151) in the spin-orbit, and the spin(1)-spin(2) case, since in isotropic Schwarzschild coordinates it vanishes identically, see (96), (97), (155), (156), and (160).

There are two possible further checks which use different approaches. As mentioned in [84] a comparison of the effective field theory NNLO spin(1)-spin(2) potential in [82] to our NNLO spin(1)-spin(2) Hamiltonian would be a very strong check of both results since the EFT results are completely independent from the ADM formalism. Also a confirming check would be the derivation of both NNLO Hamiltonians using the spin-precession method shown in [51]. Due to the complicated structure of these comparisons they will be postponed to later publications. A very recent check of the NNLO spin-orbit Hamiltonian and the resulting equations of motion was performed in [85, 86] in harmonic gauge. Furthermore in [86] the near-zone metric was determined which is an important step towards the mentioned template calculations.

10 Conclusions and Outlook

We have derived the next-to-next-to-leading order spin-orbit and spin(1)-spin(2) Hamiltonians for binary systems. The spin-orbit Hamiltonian completes the knowledge of binary black hole dynamics up to and including 3.5PN order if the objects are rapidly rotating. For neutron stars also the leading order cubic-in-spin Hamiltonians are needed as the results in [60, 61] are valid for black holes only and tidal deformation effects become very important [71–73]. The Hamiltonians were checked using two methods. The fulfillment of the global approximate Poincaré algebra was a major criterion for the correctness of the derived Hamiltonians in the extended ADM formalism. During this check the center-of-mass vectors could be determined uniquely from an ansatz. Since the approximate Poincaré algebra is not sensitive to the static part of the spin(1)-spin(2) Hamiltonian and fixed only the difference of the two coefficients at the highest order in G of the spin-orbit Hamiltonian we performed further checks. The most simple test is a linear-in-spin approximation of the Hamiltonian of a test-spin moving near a stationary Kerr black hole. We rederived the test-spin Hamiltonian from [190] in a different manner in Sect. 9 (avoiding the use of Dirac brackets). A comparison was straightforward as the same gauge was used. A more elaborate test is the recalculation of both NNLO Hamiltonians via the spin-precession frequency method in [51] and will be part of a further publication. Also a comparison of the NNLO spin(1)-spin(2) Hamiltonian with the NNLO spin(1)-spin(2) potential given in [82] will be part of a further publication and would be a very strong check, because the derivation of this potential is completely independent from the ADM formalism. The most important

confirmation to date is the independent derivation of the NNLO spin-orbit acceleration in harmonic gauge [85].

The results given in this article complete the knowledge of the post-Newtonian approximate dynamics for binary black holes up to 3.5PN. For general compact objects like neutron stars the leading order cubic-in-spin Hamiltonians are still unknown. The NNLO spin(1)-spin(2) Hamiltonian is at 4PN, if both objects are rapidly rotating, but there are still some tasks left to get the full post-Newtonian approximate dynamics up to and including 4PN. For general compact objects the leading order quartic-in-spin Hamiltonians are also unknown, they are only known for black holes.⁷ Furthermore the NNLO spin(1)² Hamiltonian – which is also at 4PN if the object is maximally rotating and maybe stronger than NNLO spin(1)-spin(2) – is completely unknown. Last but not least the G^3 up to the G^5 corrections to the 4PN point-mass Hamiltonian are also still unknown [191].

To get reasonable results for the templates the far-zone radiation field also must be calculated at higher order in the post-Newtonian approximation and also at higher orders in spin. The energy and angular momentum loss are also not known at a post-Newtonian order corresponding to next-to-next-to-leading order linear-in-spin. If radiation and fluxes are known at such high orders also a parameterization is necessary. These three major ingredients are needed for the analytical description of gravitational wave templates which are very sensitive to higher order post-Newtonian and spin corrections. Analytical results are still important because for spinning binaries the parameter space (masses and spin-directions of the components) is very large and numerical simulations are so time consuming that they cannot be used to cover the whole parameter space.⁸

Acknowledgments This work is supported by the Deutsche Forschungsgemeinschaft (DFG) through projects GRK 1523 “Quanten- und Gravitationsfelder,” STE 2017/1-1, and SFB/TR7. We gratefully acknowledge P. Jaranowski for sharing insight in 3PN point-mass calculations, for useful discussions on integration procedures, for providing several testintegrals, and for useful discussions on calculating the spin-precession frequency. In particular we thank T. Damour for very useful comments and hints regarding the calculation of d -dimensional short-range behavior of certain integrals. Many useful suggestions by an anonymous referee are gratefully acknowledged. We additionally thank M. Levi for pleasant collaboration on the comparison of our result with the potential obtained within the effective field theory approach. We also thank S. Hergt and M. Tessmer for many useful discussions about the Poincaré algebra and orbital parameterization issues, respectively. We further gratefully acknowledge useful discussions with D. Brizuela on xPERT and doing perturbation theory in arbitrary dimensions, with M. Q. Huber on three-body integral related Appell F_4 functions, with H. Witek on numerical relativity and literature on some special topics, and with T. J. Rothe on implementation issues in MATHEMATICA and for giving us some hints to mathematical theorems.

References

- [1] A. Einstein, “Zur allgemeinen Relativitätstheorie,” *Sitz.-Ber. Preuß. Akad. Wiss.* **XLIV** (1915) 778–785.
- [2] A. Einstein, “Zur allgemeinen Relativitätstheorie (Nachtrag),” *Sitz.-Ber. Preuß. Akad. Wiss.* **XLVI** (1915) 799–801.
- [3] A. Einstein, “Erklärung der Perihelbewegung des Merkur aus der allgemeinen Relativitätstheorie,” *Sitz.-Ber. Preuß. Akad. Wiss.* **XLVII** (1915) 831–839.
- [4] A. Einstein, “Die Feldgleichungen der Gravitation,” *Sitz.-Ber. Preuß. Akad. Wiss.* **XLVIII** (1915) 844–847.
- [5] A. Einstein, “Die Grundlage der allgemeinen Relativitätstheorie,” *Ann. Phys. (Berlin)* **354** (1916) 769–822.

⁷In [74] the authors argued that the spin(1)⁴ Hamiltonians derived in [61] are incomplete.

⁸In [192] they estimate that the simulation of non-spinning binaries for eight orbits for mass ratio 1:1 consumes ca. 200 000 and for 1:10 up to two million CPU hours.

- [6] H. Stephani, D. Kramer, M. A. H. MacCallum, C. Hoenselaers, and E. Herlt, *Exact Solutions of Einstein's Field Equations*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, The Edinburgh Building, Cambridge CB2 2RU, UK, 2nd ed., 2003.
- [7] J. B. Griffiths and J. Podolský, *Exact Space-Times in Einstein's General Relativity*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, The Edinburgh Building, Cambridge CB2 2RU, UK, 2009.
- [8] K. Schwarzschild, "Über das Gravitationsfeld eines Massenpunktes nach der Einsteinschen Theorie," *Sitz.-Ber. Preuß. Akad. Wiss.* (1916) 189–196.
- [9] R. P. Kerr, "Gravitational field of a spinning mass as an example of algebraically special metrics," *Phys. Rev. Lett.* **11** (1963) 237–238.
- [10] C. Cutler, T. A. Apostolatos, L. Bildsten, L. S. Finn, É. E. Flanagan, D. Kennefick, D. M. Markovic, A. Ori, E. Poisson, G. J. Sussman, and K. S. Thorne, "The last three minutes: Issues in gravitational-wave measurements of coalescing compact binaries," *Phys. Rev. Lett.* **70** (1993) 2984–2987, [arXiv:astro-ph/9208005](https://arxiv.org/abs/astro-ph/9208005).
- [11] C. Reisswig, S. Husa, L. Rezzolla, E. N. Dorband, D. Pollney, and J. Seiler, "Gravitational-wave detectability of equal-mass black-hole binaries with aligned spins," *Phys. Rev. D* **80** (2009) 124026, [arXiv:0907.0462 \[gr-qc\]](https://arxiv.org/abs/gr-qc/0907.0462).
- [12] "GEO600 home page." Electronic adress, May, 2011. <http://www.geo600.org>.
- [13] "VIRGO collaboration." Electronic adress, Oct., 2011. <https://www.cascina.virgo.infn.it/>.
- [14] "advanced LIGO home page." Electronic adress, May, 2011. <http://advancedligo.mit.edu>.
- [15] R. A. Hulse and J. H. Taylor, "Discovery of a pulsar in a binary system," *ApJ* **195** (1975) L51–L53.
- [16] M. Kramer, I. H. Stairs, R. N. Manchester, M. A. McLaughlin, A. G. Lyne, R. D. Ferdman, M. Burgay, D. R. Lorimer, A. Possenti, N. D'Amico, J. M. Sarkissian, G. B. Hobbs, J. E. Reynolds, P. C. C. Freire, and F. Camilo, "Tests of general relativity from timing the double pulsar," *Science* **314** (2006) 97–102, [arXiv:astro-ph/0609417](https://arxiv.org/abs/astro-ph/0609417).
- [17] M. Kramer and N. Wex, "The double pulsar system: a unique laboratory for gravity," *Class. Quant. Grav.* **26** (2009) 073001.
- [18] J. J. Hermes, M. Kilic, W. R. Brown, D. E. Winget, C. A. Prieto, A. Gianninas, A. S. Mukadam, A. Cabrera-Lavers, and S. J. Kenyon, "Rapid orbital decay in the 12.75-minute WD+WD binary J0651+2844," [arXiv:1208.5051 \[astro-ph.SR\]](https://arxiv.org/abs/1208.5051).
- [19] G. Schäfer, "The gravitational quadrupole radiation-reaction force and the canonical formalism of ADM," *Ann. Phys. (N.Y.)* **161** (1985) 81–100.
- [20] H. A. Lorentz and J. Droste, "De beweging van een stelsel lichamen onder den invloed van hunne onderlinge aantrekking, behandeld volgens de theorie van Einstein, I," *Versl. K. Akad. Wet. (Amsterdam)* **26** (1917) 392.
H. A. Lorentz and J. Droste, "De beweging van een stelsel lichamen onder den invloed van hunne onderlinge aantrekking, behandeld volgens de theorie van Einstein, II," *Versl. K. Akad. Wet. (Amsterdam)* **26** (1917) 649.
- [21] A. Einstein, L. Infeld, and B. Hoffmann, "The gravitational equations and the problem of motion," *Ann. Math.* **39** (1938) 65–100.
- [22] R. L. Arnowitt, S. Deser, and C. W. Misner, "The dynamics of general relativity," in *Gravitation: An Introduction to Current Research*, L. Witten, ed., pp. 227–265. John Wiley, New York, 1962.
R. L. Arnowitt, S. Deser, and C. W. Misner, "Republication of: The dynamics of general relativity," *Gen. Relativ. Gravit.* **40** (2008) 1997–2027, [arXiv:gr-qc/0405109](https://arxiv.org/abs/gr-qc/0405109).
- [23] T. Ohta, H. Okamura, T. Kimura, and K. Hiida, "Coordinate condition and higher order gravitational potential in canonical formalism," *Prog. Theor. Phys.* **51** (1974) 1598–1612.
- [24] T. Damour and G. Schäfer, "Lagrangians for n point masses at the second post-Newtonian approximation of general relativity," *Gen. Relativ. Gravit.* **17** (1985) 879–905.
- [25] T. Damour and G. Schäfer, "Higher-order relativistic periastron advances and binary pulsars," *Nuovo Cim. B* **101** (1988) 127–176.
- [26] G. Schäfer, "The ADM Hamiltonian at the postlinear approximation," *Gen. Relativ. Gravit.* **18** (1986) 255–270.

- [27] P. Jaranowski and G. Schäfer, “Third post-Newtonian higher order ADM Hamilton dynamics for two-body point-mass systems,” *Phys. Rev. D* **57** (1998) 7274–7291, [arXiv:gr-qc/9712075](#).
- [28] T. Kimura and T. Toiya, “Potential in the canonical formalism of gravity,” *Prog. Theor. Phys.* **48** (1972) 316–328.
- [29] P. Jaranowski and G. Schäfer, “Binary black-hole problem at the third post-Newtonian approximation in the orbital motion: Static part,” *Phys. Rev. D* **60** (1999) 124003, [arXiv:gr-qc/9906092](#).
- [30] T. Damour, P. Jaranowski, and G. Schäfer, “Poincaré invariance in the ADM Hamiltonian approach to the general relativistic two-body problem,” *Phys. Rev. D* **62** (2000) 021501(R), [arXiv:gr-qc/0003051](#).
- [31] T. Damour, P. Jaranowski, and G. Schäfer, “Dimensional regularization of the gravitational interaction of point masses,” *Phys. Lett. B* **513** (2001) 147–155, [arXiv:gr-qc/0105038](#).
- [32] P. Jaranowski and G. Schäfer, “Radiative 3.5 post-Newtonian ADM Hamiltonian for many-body point-mass systems,” *Phys. Rev. D* **55** (1997) 4712–4722.
- [33] C. Königsdörffer, G. Faye, and G. Schäfer, “The binary black-hole dynamics at the third-and-a-half post-Newtonian order in the ADM-formalism,” *Phys. Rev. D* **68** (2003) 044004, [arXiv:gr-qc/0305048](#).
- [34] L. Blanchet, “Gravitational radiation from post-Newtonian sources and inspiralling compact binaries,” *Living Rev. Relativity* **9** (2006) 4. <http://www.livingreviews.org/lrr-2006-4>.
- [35] T. Futamase and Y. Itoh, “The post-Newtonian approximation for relativistic compact binaries,” *Living Rev. Relativity* **10** (2007) 2. <http://www.livingreviews.org/lrr-2007-2>.
- [36] M. E. Pati and C. M. Will, “Post-Newtonian gravitational radiation and equations of motion via direct integration of the relaxed Einstein equations: Foundations,” *Phys. Rev. D* **62** (2000) 124015, [arXiv:gr-qc/0007087](#).
- [37] T. Damour and G. Esposito-Farese, “Testing gravity to second post-Newtonian order: A field theory approach,” *Phys. Rev. D* **53** (1996) 5541–5578, [arXiv:gr-qc/9506063](#) [gr-qc].
- [38] W. D. Goldberger and I. Z. Rothstein, “An effective field theory of gravity for extended objects,” *Phys. Rev. D* **73** (2006) 104029, [arXiv:hep-th/0409156](#).
- [39] J. B. Gilmore and A. Ross, “Effective field theory calculation of second post-Newtonian binary dynamics,” *Phys. Rev. D* **78** (2008) 124021, [arXiv:0810.1328](#) [gr-qc].
- [40] B. Kol and M. Smolkin, “Dressing the post-Newtonian two-body problem and classical effective field theory,” *Phys. Rev. D* **80** (2009) 124044, [arXiv:0910.5222](#) [hep-th].
- [41] S. Foffa and R. Sturani, “Effective field theory calculation of conservative binary dynamics at third post-Newtonian order,” *Phys. Rev. D* **84** (2011) 044031, [arXiv:1104.1122](#) [gr-qc].
- [42] S. Foffa and R. Sturani, “The dynamics of the gravitational two-body problem in the post-Newtonian approximation at quadratic order in the Newton’s constant,” [arXiv:1206.7087](#) [gr-qc].
- [43] B. M. Barker and R. F. O’Connell, “Gravitational two-body problem with arbitrary masses, spins, and quadrupole moments,” *Phys. Rev. D* **12** (1975) 329–335.
- [44] P. D. D’Eath, “Interaction of two black holes in the slow-motion limit,” *Phys. Rev. D* **12** (1975) 2183–2199.
- [45] B. M. Barker and R. F. O’Connell, “The gravitational interaction: Spin, rotation, and quantum effects—a review,” *Gen. Relativ. Gravit.* **11** (1979) 149–175.
- [46] K. S. Thorne and J. B. Hartle, “Laws of motion and precession for black holes and other bodies,” *Phys. Rev. D* **31** (1985) 1815–1837.
- [47] H. Goenner, U. Gralewski, and K. Westpfahl, “Gravitative Selbstkräfte und Strahlungsverluste klassischer Spinteilchen (erste Näherung),” *Z. Phys.* **207** (1967) 186–208.
- [48] F. Bennewitz and K. Westpfahl, “Selbstwechselwirkung von Gravitationsfeldern schnell bewegter Pol-Dipolquellen,” *Commun. math. Phys.* **23** (1971) 296–318.
- [49] H. Tagoshi, A. Ohashi, and B. J. Owen, “Gravitational field and equations of motion of spinning compact binaries to 2.5 post-Newtonian order,” *Phys. Rev. D* **63** (2001) 044006, [arXiv:gr-qc/0010014](#).
- [50] G. Faye, L. Blanchet, and A. Buonanno, “Higher-order spin effects in the dynamics of compact

- binaries. I. Equations of motion,” *Phys. Rev. D* **74** (2006) 104033, [arXiv:gr-qc/0605139](#).
- [51] T. Damour, P. Jaranowski, and G. Schäfer, “Hamiltonian of two spinning compact bodies with next-to-leading order gravitational spin-orbit coupling,” *Phys. Rev. D* **77** (2008) 064032, [arXiv:0711.1048 \[gr-qc\]](#).
- [52] J. Steinhoff, S. Hergt, and G. Schäfer, “Next-to-leading order gravitational spin(1)-spin(2) dynamics in Hamiltonian form,” *Phys. Rev. D* **77** (2008) 081501(R), [arXiv:0712.1716 \[gr-qc\]](#).
- [53] J. Steinhoff, S. Hergt, and G. Schäfer, “Spin-squared Hamiltonian of next-to-leading order gravitational interaction,” *Phys. Rev. D* **78** (2008) 101503(R), [arXiv:0809.2200 \[gr-qc\]](#).
- [54] D. L. Perrodin, “Subleading spin-orbit correction to the Newtonian potential in effective field theory formalism,” in *Proceedings of the 12th Marcel Grossmann Meeting on General Relativity*. World Scientific, Singapore, 2011. [arXiv:1005.0634 \[gr-qc\]](#).
- [55] R. A. Porto, “Next to leading order spin-orbit effects in the motion of inspiralling compact binaries,” *Class. Quant. Grav.* **27** (2010) 205001, [arXiv:1005.5730 \[gr-qc\]](#).
- [56] M. Levi, “Next-to-leading order gravitational spin-orbit coupling in an effective field theory approach,” *Phys. Rev. D* **82** (2010) 104004, [arXiv:1006.4139 \[gr-qc\]](#).
- [57] R. A. Porto and I. Z. Rothstein, “Spin(1)spin(2) effects in the motion of inspiralling compact binaries at third order in the post-Newtonian expansion,” *Phys. Rev. D* **78** (2008) 044012, [arXiv:0802.0720 \[gr-qc\]](#).
R. A. Porto and I. Z. Rothstein, “Erratum: Spin(1)spin(2) effects in the motion of inspiralling compact binaries at third order in the post-Newtonian expansion,” *Phys. Rev. D* **81** (2010) 029904(E).
- [58] M. Levi, “Next-to-leading order gravitational spin1-spin2 coupling with Kaluza-Klein reduction,” *Phys. Rev. D* **82** (2010) 064029, [arXiv:0802.1508 \[gr-qc\]](#).
- [59] H. Wang, J. Steinhoff, J. Zeng, and G. Schäfer, “Leading-order spin-orbit and spin(1)-spin(2) radiation-reaction Hamiltonians,” *Phys. Rev. D* **84** (2011) 124005, [arXiv:1109.1182 \[gr-qc\]](#).
- [60] S. Hergt and G. Schäfer, “Higher-order-in-spin interaction Hamiltonians for binary black holes from source terms of Kerr geometry in approximate ADM coordinates,” *Phys. Rev. D* **77** (2008) 104001, [arXiv:0712.1515 \[gr-qc\]](#).
- [61] S. Hergt and G. Schäfer, “Higher-order-in-spin interaction Hamiltonians for binary black holes from Poincaré invariance,” *Phys. Rev. D* **78** (2008) 124004, [arXiv:0809.2208 \[gr-qc\]](#).
- [62] E. Poisson, “Gravitational waves from inspiraling compact binaries: The quadrupole-moment term,” *Phys. Rev. D* **57** (1998) 5287–5290, [arXiv:gr-qc/9709032](#).
- [63] R. A. Porto and I. Z. Rothstein, “Next to leading order spin(1)spin(1) effects in the motion of inspiralling compact binaries,” *Phys. Rev. D* **78** (2008) 044013, [arXiv:0804.0260 \[gr-qc\]](#).
R. A. Porto and I. Z. Rothstein, “Erratum: Next to leading order spin(1)spin(1) effects in the motion of inspiralling compact binaries,” *Phys. Rev. D* **81** (2010) 029905(E).
- [64] J. Steinhoff and G. Schäfer, “Comment on two recent papers regarding next-to-leading order spin-spin effects in gravitational interaction,” *Phys. Rev. D* **80** (2009) 088501, [arXiv:0903.4772 \[gr-qc\]](#).
- [65] S. Hergt, J. Steinhoff, and G. Schäfer, “The reduced Hamiltonian for next-to-leading-order spin-squared dynamics of general compact binaries,” *Class. Quant. Grav.* **27** (2010) 135007, [arXiv:1002.2093 \[gr-qc\]](#).
- [66] G. Schäfer, “Three-body Hamiltonian in General Relativity,” *Phys. Lett. A* **123** (1987) 336–339.
- [67] T. Ohta, T. Kimura, and K. Hiida, “Can the effect of distant matter on physical observables be observed?,” *Nuovo Cim. B* **27** (1975) 103–120.
- [68] C. O. Lousto and H. Nakano, “Three-body equations of motion in successive post-Newtonian approximations,” *Class. Quant. Grav.* **25** (2008) 195019, [arXiv:0710.5542 \[gr-qc\]](#).
- [69] Y.-Z. Chu, “n-body problem in general relativity up to the second post-Newtonian order from perturbative field theory,” *Phys. Rev. D* **79** (2009) 044031, [arXiv:0812.0012 \[gr-qc\]](#).
- [70] J. Hartung and J. Steinhoff, “Next-to-leading order spin-orbit and spin(a)-spin(b) Hamiltonians for n gravitating spinning compact objects,” *Phys. Rev. D* **83** (2011) 044008, [arXiv:1011.1179 \[gr-qc\]](#).

- [71] T. Damour and A. Nagar, “Effective one body description of tidal effects in inspiralling compact binaries,” *Phys. Rev. D* **81** (2010) 084016, [arXiv:0911.5041 \[gr-qc\]](#).
- [72] J. E. Vines and É. E. Flanagan, “Post-1-Newtonian quadrupole tidal interactions in binary systems,” [arXiv:1009.4919 \[gr-qc\]](#).
- [73] D. Bini, T. Damour, and G. Faye, “Effective action approach to higher-order relativistic tidal interactions in binary systems and their effective one body description,” *Phys. Rev. D* **85** (2012) 124034, [arXiv:1202.3565 \[gr-qc\]](#).
- [74] J. Steinhoff and D. Puetzfeld, “Influence of internal structure on the motion of test bodies in extreme mass ratio situations,” *Phys. Rev. D* **86** (2012) 044033, [arXiv:1205.3926 \[gr-qc\]](#).
- [75] J. Steinhoff and G. Schäfer, “Canonical formulation of self-gravitating spinning-object systems,” *Europhys. Lett.* **87** (2009) 50004, [arXiv:0907.1967 \[gr-qc\]](#).
- [76] J. Steinhoff, “Canonical formulation of spin in general relativity,” *Ann. Phys. (Berlin)* **523** (2011) 296–353, [arXiv:1106.4203 \[gr-qc\]](#).
- [77] J. Steinhoff and H. Wang, “Canonical formulation of gravitating spinning objects at 3.5 post-Newtonian order,” *Phys. Rev. D* **81** (2010) 024022, [arXiv:0910.1008 \[gr-qc\]](#).
- [78] J. Steinhoff, G. Schäfer, and S. Hergt, “ADM canonical formalism for gravitating spinning objects,” *Phys. Rev. D* **77** (2008) 104018, [arXiv:0805.3136 \[gr-qc\]](#).
- [79] J. Hartung and J. Steinhoff, “Next-to-next-to-leading order post-Newtonian spin-orbit Hamiltonian for self-gravitating binaries,” *Ann. Phys. (Berlin)* **523** (2011) 783–790, [arXiv:1104.3079 \[gr-qc\]](#).
- [80] J. Hartung and J. Steinhoff, “Next-to-next-to-leading order post-Newtonian spin(1)-spin(2) Hamiltonian for self-gravitating binaries,” *Ann. Phys. (Berlin)* **523** (2011) 919–924, [arXiv:1107.4294 \[gr-qc\]](#).
- [81] J. Hartung, *Binärsysteme kompakter Objekte in hoher postnewtonscher Näherung in den Eigendrehimpuls-Wechselwirkungen*. PhD thesis, Friedrich-Schiller-Universität Jena, 2012.
- [82] M. Levi, “Binary dynamics from spin1-spin2 coupling at fourth post-Newtonian order,” *Phys. Rev. D* **85** (2012) 064043, [arXiv:1107.4322 \[gr-qc\]](#).
- [83] S. Hergt, *Hamiltonsche Formulierung und Behandlung nichtlinearer Eigendrehimpulsbeiträge in Binärsystemen der Allgemeinen Relativitätstheorie*. PhD thesis, Friedrich-Schiller-Universität Jena, 2011.
- [84] S. Hergt, J. Steinhoff, and G. Schäfer, “Elimination of the spin supplementary condition in the effective field theory approach to the post-Newtonian approximation,” *Ann. Phys. (N.Y.)* **327** (2012) 1494–1537, [arXiv:1110.2094 \[gr-qc\]](#).
- [85] S. Marsat, A. Bohe, G. Faye, and L. Blanchet, “Next-to-next-to-leading order spin-orbit effects in the equations of motion of compact binary systems,” *Class. Quant. Grav.* **30** (2013) 055007, [arXiv:1210.4143 \[gr-qc\]](#).
- [86] A. Bohe, S. Marsat, G. Faye, and L. Blanchet, “Next-to-next-to-leading order spin-orbit effects in the near-zone metric and precession equations of compact binaries,” [arXiv:1212.5520 \[gr-qc\]](#).
- [87] T. Ledvinka, G. Schäfer, and J. Bičák, “Relativistic closed-form Hamiltonian for many-body gravitating systems in the post-Minkowskian approximation,” *Phys. Rev. Lett.* **100** (2008) 251101, [arXiv:0807.0214 \[gr-qc\]](#).
- [88] M. Tessmer, *Motion and gravitational wave emission of spinning compact binaries*. PhD thesis, Friedrich-Schiller-Universität Jena, 2011.
- [89] T. Damour and N. Deruelle, “General relativistic celestial mechanics of binary systems. I. the post-Newtonian motion,” *Ann. Inst. H. Poincaré A* **43** (1985) 107–132. http://www.numdam.org/item?id=AIHPA_1985__43_1_107_0.
- [90] G. Schäfer and N. Wex, “Second post-Newtonian motion of compact binaries,” *Phys. Lett. A* **174** (1993) 196–205.
G. Schäfer and N. Wex, “Erratum: Second post-Newtonian motion of compact binaries,” *Phys. Lett. A* **177** (1993) 461(E).
- [91] R.-M. Memmesheimer, A. Gopakumar, and G. Schäfer, “Third post-Newtonian accurate generalized quasi-Keplerian parametrization for compact binaries in eccentric orbits,” *Phys. Rev. D* **70** (2004) 104011, [arXiv:gr-qc/0407049](#).

- [92] N. Wex, “The second post-Newtonian motion of compact binary-star systems with spin,” *Class. Quant. Grav.* **12** (1995) 983–1005.
- [93] C. Königsdörffer and A. Gopakumar, “Post-Newtonian accurate parametric solution to the dynamics of spinning compact binaries in eccentric orbits: The leading order spin-orbit interaction,” *Phys. Rev. D* **71** (2005) 024039, [arXiv:gr-qc/0501011](#).
- [94] C. Königsdörffer and A. Gopakumar, “Phasing of gravitational waves from inspiralling eccentric binaries at the third-and-a-half post-Newtonian order,” *Phys. Rev. D* **73** (2006) 124012, [arXiv:gr-qc/0603056](#).
- [95] M. Tessmer, “Gravitational waveforms from unequal-mass binaries with arbitrary spins under leading order spin-orbit coupling,” *Phys. Rev. D* **80** (2009) 124034, [arXiv:0910.5931 \[gr-qc\]](#).
- [96] M. Tessmer, J. Hartung, and G. Schäfer, “Motion and gravitational wave forms of eccentric compact binaries with orbital-angular-momentum-aligned spins under next-to-leading order in spin-orbit and leading order in spin(1)–spin(2) and spin-squared couplings,” *Class. Quant. Grav.* **27** (2010) 165005, [arXiv:1003.2735 \[gr-qc\]](#).
- [97] M. Tessmer, J. Hartung, and G. Schäfer, “Aligned spins: Orbital elements, decaying orbits, and last stable circular orbit to high post-Newtonian orders,” *Class. Quant. Grav.* **30** (2013) 015007, [arXiv:1207.6961 \[gr-qc\]](#).
- [98] W. B. Bonnor, “Spherical gravitational waves,” *Phil. Trans. R. Soc. A* **251** (1959) 233–271.
- [99] K. S. Thorne, “Multipole expansions of gravitational radiation,” *Rev. Mod. Phys.* **52** (1980) 299–339.
- [100] L. Blanchet and T. Damour, “Radiative gravitational fields in general relativity I. General structure of the field outside the source,” *Phil. Trans. R. Soc. A* **320** (1986) 379–430. <http://www.jstor.org/stable/37878>.
- [101] L. Blanchet and G. Schäfer, “Higher order gravitational radiation losses in binary systems,” *Mon. Not. R. Astron. Soc.* **239** (1989) 845–867. <http://adsabs.harvard.edu/abs/1989MNRAS.239..845B>.
L. Blanchet and G. Schäfer, “Higher order gravitational radiation losses in binary systems: Erratum,” *Mon. Not. R. Astron. Soc.* **242** (1990) 704.
- [102] L. E. Kidder, “Coalescing binary systems of compact objects to (post)^{5/2}-Newtonian order. V. Spin effects,” *Phys. Rev. D* **52** (1995) 821–847, [arXiv:gr-qc/9506022](#).
- [103] L. Blanchet, A. Buonanno, and G. Faye, “Higher-order spin effects in the dynamics of compact binaries. II. Radiation field,” *Phys. Rev. D* **74** (2006) 104034, [arXiv:gr-qc/0605140](#).
L. Blanchet, A. Buonanno, and G. Faye, “Erratum: Higher-order spin effects in the dynamics of compact binaries. II. Radiation field,” *Phys. Rev. D* **75** (2007) 049903(E).
L. Blanchet, A. Buonanno, and G. Faye, “Erratum: Higher-order spin effects in the dynamics of compact binaries. II. Radiation field,” *Phys. Rev. D* **81** (2010) 089901(E).
- [104] A. Buonanno, G. Faye, and T. Hinderer, “Spin effects on gravitational waves from inspiraling compact binaries at second post-Newtonian order,” [arXiv:1209.6349 \[gr-qc\]](#).
- [105] L. Blanchet, A. Buonanno, and G. Faye, “Tail-induced spin-orbit effect in the gravitational radiation of compact binaries,” *Phys. Rev. D* **84** (2011) 064041, [arXiv:1104.5659 \[gr-qc\]](#).
- [106] R. A. Porto, A. Ross, and I. Z. Rothstein, “Spin induced multipole moments for the gravitational wave amplitude from binary inspirals to 2.5 post-Newtonian order,” *JCAP* **1209** (2012) 028, [arXiv:1203.2962 \[gr-qc\]](#).
- [107] R. A. Porto, A. Ross, and I. Z. Rothstein, “Spin induced multipole moments for the gravitational wave flux from binary inspirals to third post-Newtonian order,” *JCAP* **1103** (2011) 009, [arXiv:1007.1312 \[gr-qc\]](#).
- [108] M. Turner and C. M. Will, “Post-Newtonian gravitation bremsstrahlung,” *ApJ* **220** (1978) 1107.
- [109] D. V. Gal’tsov, A. A. Matiukhin, and V. I. Petukhov, “Relativistic corrections to the gravitational radiation of a binary system and the fine structure of the spectrum,” *Phys. Lett. A* **77** (1980) 387.
- [110] V. Pierro, I. M. Pinto, E. Spallicci, E. Laserra, and F. Recano, “Fast and accurate computational tools for gravitational waveforms from binary stars with any orbital eccentricity,” *Mon. Not. R. Astron. Soc.* **325** (2001) 358–372.
- [111] W. Junker and G. Schäfer, “Binary systems - higher order gravitational radiation damping and wave

- emission,” *Mon. Not. R. Astron. Soc.* **254** (1992) 146–164.
<http://adsabs.harvard.edu/abs/1992MNRAS.254...146J>.
- [112] C. Moreno-Garrido, J. Buitrago, and E. Mediavilla, “Spectral analysis of the gravitational radiation emitted by binary systems in moderately eccentric orbits - application to coalescing binaries,” *Mon. Not. R. Astron. Soc.* **266** (1994) 16.
<http://adsabs.harvard.edu/abs/1994MNRAS.266...16M>.
- [113] C. Moreno-Garrido, E. Mediavilla, and J. Buitrago, “Gravitational radiation from point masses in elliptical orbits: spectral analysis and orbital parameters,” *Mon. Not. R. Astron. Soc.* **274** (1995) 115–126. <http://adsabs.harvard.edu/abs/1995MNRAS.274...115M>.
- [114] A. Gopakumar and B. R. Iyer, “Second post-Newtonian gravitational wave polarizations for compact binaries in elliptical orbits,” *Phys. Rev. D* **65** (2002) 084011, [arXiv:0110100 \[gr-qc\]](https://arxiv.org/abs/gr-qc/0110100).
- [115] M. Tessmer and A. Gopakumar, “Accurate and efficient gravitational waveforms for certain galactic compact binaries,” *Mon. Not. R. Astron. Soc.* **374** (2007) 721–728, [arXiv:gr-qc/0610139](https://arxiv.org/abs/gr-qc/0610139).
- [116] M. Tessmer and G. Schäfer, “Full-analytic frequency-domain 1pN-accurate gravitational wave forms from eccentric compact binaries,” *Phys. Rev. D* **82** (2010) 124064, [arXiv:1006.3714v2 \[gr-qc\]](https://arxiv.org/abs/1006.3714v2).
- [117] M. Tessmer and G. Schäfer, “Full-analytic frequency-domain gravitational wave forms from eccentric compact binaries to 2PN accuracy,” *Ann. Phys. (Berlin)* **523** (2011) 813–864, [arXiv:1012.3894 \[gr-qc\]](https://arxiv.org/abs/1012.3894).
- [118] A. Gopakumar and B. R. Iyer, “Gravitational waves from inspiraling compact binaries: Angular momentum flux, evolution of the orbital elements, and the waveform to the second post-Newtonian order,” *Phys. Rev. D* **56** (1997) 7708–7731, [arXiv:gr-qc/0110100](https://arxiv.org/abs/gr-qc/0110100).
- [119] T. Damour, A. Gopakumar, and B. R. Iyer, “Phasing of gravitational waves from inspiralling eccentric binaries,” *Phys. Rev. D* **70** (2004) 064028, [arXiv:gr-qc/0404128](https://arxiv.org/abs/gr-qc/0404128).
- [120] S. Chandrasekhar and F. P. Esposito, “The 2 1/2-Post-Newtonian Equations of Hydrodynamics and Radiation Reaction in General Relativity,” *ApJ* **160** (1970) 153.
- [121] L. S. Finn, “Detection, measurement, and gravitational radiation,” *Phys. Rev. D* **46** (1992) 5236–5249, [arXiv:gr-qc/9609027](https://arxiv.org/abs/gr-qc/9609027).
- [122] T. A. Apostolatos, “Search templates for gravitational waves from precessing, inspiraling binaries,” *Phys. Rev. D* **52** (1995) 605–620.
- [123] T. Damour, B. R. Iyer, and B. S. Sathyaprakash, “Frequency-domain p-approximant filters for time-truncated inspiral gravitational wave signals from compact binaries,” *Phys. Rev. D* **62** (2000) 084036, [arXiv:gr-qc/0001023](https://arxiv.org/abs/gr-qc/0001023).
- [124] T. Damour, B. R. Iyer, and B. S. Sathyaprakash, “Comparison of search templates for gravitational waves from binary inspiral,” *Phys. Rev. D* **63** (2001) 044023, [arXiv:gr-qc/0010009](https://arxiv.org/abs/gr-qc/0010009).
T. Damour, B. R. Iyer, and B. S. Sathyaprakash, “Erratum: Comparison of search templates for gravitational waves from binary inspiral,” *Phys. Rev. D* **72** (2005) 029902(E).
- [125] P. Ajith, S. Babak, Y. Chen, M. Hewitson, B. Krishnan, A. M. Sintes, J. T. Whelan, B. Brügmann, P. Diener, N. Dorband, J. Gonzalez, M. Hannam, S. Husa, D. Pollney, L. Rezzolla, L. Santamaría, U. Sperhake, and J. Thornburg, “Template bank for gravitational waveforms from coalescing binary black holes: Nonspinning binaries,” *Phys. Rev. D* **77** (2008) 104017, [arXiv:0710.2335 \[gr-qc\]](https://arxiv.org/abs/0710.2335).
P. Ajith, S. Babak, Y. Chen, M. Hewitson, B. Krishnan, A. M. Sintes, J. T. Whelan, B. Brügmann, P. Diener, N. Dorband, J. Gonzalez, M. Hannam, S. Husa, D. Pollney, L. Rezzolla, L. Santamaría, U. Sperhake, and J. Thornburg, “Erratum: Template bank for gravitational waveforms from coalescing binary black holes: Nonspinning binaries,” *Phys. Rev. D* **79** (2009) 129901(E).
- [126] A. Buonanno, Y. Chen, and M. Vallisneri, “Detecting gravitational waves from precessing binaries of spinning compact objects: Adiabatic limit,” *Phys. Rev. D* **67** (2003) 104025, [arXiv:gr-qc/0211087](https://arxiv.org/abs/gr-qc/0211087).
A. Buonanno, Y. Chen, and M. Vallisneri, “Erratum: Detecting gravitational waves from precessing binaries of spinning compact objects: Adiabatic limit,” *Phys. Rev. D* **74** (2006) 029904(E).
- [127] T. Damour and A. Nagar, “An improved analytical description of inspiralling and coalescing black-hole binaries,” *Phys. Rev. D* **79** (2009) 081503(R), [arXiv:0902.0136 \[gr-qc\]](https://arxiv.org/abs/0902.0136).

- [128] A. Buonanno, Y. Pan, H. P. Pfeiffer, M. A. Scheel, L. T. Buchman, and L. E. Kidder, “Effective-one-body waveforms calibrated to numerical relativity simulations: Coalescence of non-spinning, equal-mass black holes,” *Phys. Rev. D* **79** (2009) 124028, [arXiv:0902.0790 \[gr-qc\]](#).
- [129] Y. Pan, A. Buonanno, L. T. Buchman, T. Chu, L. E. Kidder, H. P. Pfeiffer, and M. A. Scheel, “Effective-one-body waveforms calibrated to numerical relativity simulations: Coalescence of nonprecessing, spinning, equal-mass black holes,” *Phys. Rev. D* **81** (2010) 084041, [arXiv:0912.3466 \[gr-qc\]](#).
- [130] T. Damour, “Coalescence of two spinning black holes: An effective one-body approach,” *Phys. Rev. D* **64** (2001) 124013, [arXiv:gr-qc/0103018](#).
- [131] E. Barausse and A. Buonanno, “Improved effective-one-body Hamiltonian for spinning black-hole binaries,” *Phys. Rev. D* **81** (2010) 084024, [arXiv:0912.3517 \[gr-qc\]](#).
- [132] A. Nagar, “Effective one-body Hamiltonian of two spinning black holes with next-to-next-to-leading order spin-orbit coupling,” *Phys. Rev. D* **84** (2011) 084028, [arXiv:1106.4349 \[gr-qc\]](#).
- [133] E. Barausse and A. Buonanno, “Extending the effective-one-body Hamiltonian of black-hole binaries to include next-to-next-to-leading spin-orbit couplings,” *Phys. Rev. D* **84** (2011) 104027, [arXiv:1107.2904 \[gr-qc\]](#).
- [134] A. Taracchini, Y. Pan, A. Buonanno, E. Barausse, M. Boyle, T. Chu, G. Lovelace, H. P. Pfeiffer, and M. A. Scheel, “A prototype effective-one-body model for non-precessing spinning inspiral-merger-ringdown waveforms,” [arXiv:1202.0790 \[gr-qc\]](#).
- [135] T. Damour, P. Jaranowski, and G. Schäfer, “Dynamical invariants for general relativistic two-body systems at the third post-Newtonian approximation,” *Phys. Rev. D* **62** (2000) 044024, [arXiv:gr-qc/9912092](#).
- [136] T. Damour, P. Jaranowski, and G. Schäfer, “Determination of the last stable orbit for circular general relativistic binaries at the third post-Newtonian approximation,” *Phys. Rev. D* **62** (2000) 084011, [arXiv:gr-qc/0005034](#).
- [137] J. Faber, “Status of neutron star–black hole and binary neutron star simulations,” *Class. Quant. Grav.* **26** (2009) 114004.
- [138] M. D. Duez, “Numerical relativity confronts compact neutron star binaries: a review and status report,” *Class. Quant. Grav.* **27** (2010) 114002, [arXiv:0912.3529 \[gr-qc\]](#).
- [139] S. Rosswog, “Compact binary mergers: an astrophysical perspective,” [arXiv:1012.0912 \[astro-ph.HE\]](#).
- [140] M. Shibata and K. Uryū, “Simulation of merging binary neutron stars in full general relativity: $\Gamma = 2$ case,” *Phys. Rev. D* **61** (2000) 064001, [arXiv:gr-qc/9911058](#).
- [141] M. Thierfelder, S. Bernuzzi, and B. Brügmann, “Numerical relativity simulations of binary neutron stars,” *Phys. Rev. D* **84** (2011) 044012, [arXiv:1104.4751 \[gr-qc\]](#).
- [142] R. Gold, S. Bernuzzi, M. Thierfelder, B. Brügmann, and F. Pretorius, “Eccentric binary neutron star mergers,” [arXiv:1109.5128 \[gr-qc\]](#).
- [143] S. Bernuzzi, A. Nagar, M. Thierfelder, and B. Brügmann, “Tidal effects in binary neutron star coalescence,” [arXiv:1205.3403 \[gr-qc\]](#).
- [144] F. Pretorius, “Evolution of binary black-hole spacetimes,” *Phys. Rev. Lett.* **95** (2005) 121101, [arXiv:gr-qc/0507014](#).
- [145] J. G. Baker, J. Centrella, D.-I. Choi, M. Koppitz, and J. van Meter, “Gravitational-wave extraction from an inspiraling configuration of merging black holes,” *Phys. Rev. Lett.* **96** (2006) 111102, [arXiv:gr-qc/0511103](#).
- [146] C. O. Lousto and Y. Zlochower, “Foundations of multiple-black-hole evolutions,” *Phys. Rev. D* **77** (2008) 024034, [arXiv:0711.1165 \[gr-qc\]](#).
- [147] M. Campanelli, C. O. Lousto, and Y. Zlochower, “Close encounters of three black holes,” *Phys. Rev. D* **77** (2008) 101501(R), [arXiv:0710.0879 \[gr-qc\]](#).
- [148] P. Galaviz, B. Brügmann, and Z. Cao, “Numerical evolution of multiple black holes with accurate initial data,” *Phys. Rev. D* **82** (2010) 024005, [arXiv:1004.1353 \[gr-qc\]](#).
- [149] P. Galaviz, “Stability and chaos of hierarchical three black hole configurations,” *Phys. Rev. D* **84** (2011) 104038, [arXiv:1108.4485 \[gr-qc\]](#).

- [150] A. Le Tiec, E. Barausse, and A. Buonanno, “Gravitational self-force correction to the binding energy of compact binary systems,” *Phys. Rev. Lett.* **108** (2012) 131103, [arXiv:1111.5609 \[gr-qc\]](#).
- [151] L. Blanchet, A. Buonanno, and A. Le Tiec, “First law of mechanics for black hole binaries with spins,” [arXiv:1211.1060 \[gr-qc\]](#).
- [152] J. M. Martín-García, *xAct: Efficient Tensor Computer Algebra*. <http://www.xact.es/>.
- [153] S. Wolfram, *The Mathematica Book*. Wolfram Media, Champaign, IL, 5th ed., 2003.
- [154] J. M. Martín-García, “xPerm: fast index canonicalization for tensor computer algebra,” *Comp. Phys. Commun.* **179** (2008) 597–603, [arXiv:0803.0862 \[cs.SC\]](#).
- [155] D. Brizuela, J. M. Martín-García, and G. A. Mena Marugán, “xPert: computer algebra for metric perturbation theory,” *Gen. Relativ. Gravit.* **41** (2009) 2415–2431, [arXiv:0807.0824 \[gr-qc\]](#).
- [156] B. S. DeWitt, “Quantum theory of gravity. I. The canonical theory,” *Phys. Rev.* **160** (1967) 1113–1148.
- [157] T. Regge and C. Teitelboim, “Role of surface integrals in the Hamiltonian formulation of general relativity,” *Ann. Phys. (N.Y.)* **88** (1974) 286–318.
- [158] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*. W. H. FREEMAN AND COMPANY, 41 Madison Avenue, New York, 21 ed., 1973.
- [159] E. Poisson, “An advanced course in general relativity.” Electronic adress, 2002. <http://www.physics.uoguelph.ca/poisson/research/agr.pdf>. Draft of lecture notes.
- [160] É. Gourgoulhon, “3+1 formalism and bases of numerical relativity,” [arXiv:gr-qc/0703035](#).
- [161] M. Mathisson, “Neue Mechanik materieller Systeme,” *Acta Phys. Pol.* **6** (1937) 163–200.
M. Mathisson, “Republication of: New mechanics of material systems,” *Gen. Relativ. Gravit.* **42** (2010) 1011–1048.
- [162] A. Papapetrou, “Spinning test-particles in general relativity. I,” *Proc. R. Soc. A* **209** (1951) 248–258.
- [163] W. M. Tulczyjew, “Motion of multipole particles in general relativity theory,” *Acta Phys. Pol.* **18** (1959) 393–409.
- [164] W. G. Dixon, “Extended bodies in general relativity: Their description and motion,” in *Proceedings of the International School of Physics Enrico Fermi LXVII*, J. Ehlers, ed., pp. 156–219. North Holland, Amsterdam, 1979.
- [165] K. Westpfahl, “Relativistische Bewegungsprobleme. I. Das freie Spinteilchen,” *Ann. Phys. (Berlin)* **475** (1967) 113–135.
- [166] K. Westpfahl, “Relativistische Bewegungsprobleme. V. Zur allgemein-relativistischen Dynamik klassischer Spinteilchen,” *Ann. Phys. (Berlin)* **477** (1969) 345–360.
- [167] H. Goenner and K. Westpfahl, “Relativistische Bewegungsprobleme. II. Der starre Rotator,” *Ann. Phys. (Berlin)* **475** (1967) 230–240.
- [168] H. Römer and K. Westpfahl, “Relativistische Bewegungsprobleme. IV. Rotator-Spinteilchen in schwachen Gravitationsfeldern,” *Ann. Phys. (Berlin)* **477** (1969) 264–276.
- [169] K. Westpfahl, “Relativistische Bewegungsprobleme. VI. Rotator-Spinteilchen und allgemeine Relativitätstheorie,” *Ann. Phys. (Berlin)* **477** (1969) 361–371.
- [170] A. J. Hanson and T. Regge, “The relativistic spherical top,” *Ann. Phys. (N.Y.)* **87** (1974) 498–566.
- [171] I. Bailey and W. Israel, “Lagrangian dynamics of spinning particles and polarized media in general relativity,” *Commun. math. Phys.* **42** (1975) 65–82.
- [172] G. Faye, P. Jaranowski, and G. Schäfer, “Skeleton approximate solution of the einstein field equations for multiple black-hole systems,” *Phys. Rev. D* **69** (2004) 124029, [arXiv:gr-qc/0311018](#).
- [173] B. C. Xanthopoulos and T. Zannias, “Einstein gravity coupled to a massless scalar field in arbitrary spacetime dimensions,” *Phys. Rev. D* **40** (1989) 2564–2567.
- [174] L. D. Landau and E. M. Lifshitz, *The classical theory of fields*, vol. 2. Pergamon Press, 1951.
- [175] J. W. York, “Role of conformal three-geometry in the dynamics of gravitation,” *Phys. Rev. Lett.* **28** (1972) 1082–1085.
- [176] G. W. Gibbons and S. W. Hawking, “Action integrals and partition functions in quantum gravity,”

- Phys. Rev. D* **15** (1977) 2752–2756.
- [177] R. M. Wald, *General Relativity*. The University of Chicago Press, Chicago 60637, 1984.
- [178] J. W. York, “Boundary terms in the action principles of general relativity,” *Found. Phys.* **16** (1986) 249–257.
- [179] J. D. Brown and J. W. York, “Quasilocal energy and conserved charges derived from the gravitational action,” *Phys. Rev. D* **47** (1993) 1407–1419, [arXiv:gr-qc/9209012](#).
- [180] S. W. Hawking and G. T. Horowitz, “The gravitational Hamiltonian, action, entropy and surface terms,” *Class. Quant. Grav.* **13** (1996) 1487–1498, [arXiv:gr-qc/9501014](#).
- [181] D. V. Gal’tsov, “Radiation reaction in various dimensions,” *Phys. Rev. D* **66** (2002) 025016, [arXiv:hep-th/0112110](#).
- [182] V. Cardoso, Ó. J. C. Dias, and J. P. S. Lemos, “Gravitational radiation in D -dimensional spacetimes,” *Phys. Rev. D* **67** (2003) 064026, [arXiv:hep-th/0212168](#).
- [183] L. Blanchet, T. Damour, G. Esposito-Farèse, and B. R. Iyer, “Dimensional regularization of the third post-Newtonian gravitational wave generation from two point masses,” *Phys. Rev. D* **71** (2005) 124004, [arXiv:gr-qc/0503044](#).
- [184] L. Blanchet, T. Damour, and G. Esposito-Farèse, “Dimensional regularization of the third post-Newtonian dynamics of point particles in harmonic coordinates,” *Phys. Rev. D* **69** (2004) 124007, [arXiv:gr-qc/0311052](#).
- [185] D. D. Iwanenko and A. A. Sokolow, *Klassische Feldtheorie*. Akademie-Verlag GmbH, Berlin, 1953.
- [186] W. D. Goldberger, “Les Houches lectures on effective field theories and gravitational radiation,” [arXiv:hep-ph/0701129](#).
- [187] C. R. Galley, “The classical mechanics of non-conservative systems,” [arXiv:1210.2745 \[gr-qc\]](#).
- [188] T. W. B. Kibble, “Canonical variables for the interacting gravitational and Dirac fields,” *J. Math. Phys.* **4** (1963) 1433–1437.
- [189] R. Beig and N. Ó Murchadha, “The Poincaré group as the symmetry group of canonical general relativity,” *Ann. Phys. (N.Y.)* **174** (1987) 463–498.
- [190] E. Barausse, É. Racine, and A. Buonanno, “Hamiltonian of a spinning test-particle in curved spacetime,” *Phys. Rev. D* **80** (2009) 104025, [arXiv:0907.4745 \[gr-qc\]](#).
- [191] P. Jaranowski and G. Schäfer, “Towards the fourth post-Newtonian Hamiltonian for two-point-mass systems,” *Phys. Rev. D* **86** (2012) 061503, [arXiv:1207.5448 \[gr-qc\]](#).
- [192] I. Hinder, “The current status of binary black hole simulations in numerical relativity,” *Class. Quant. Grav.* **27** (2010) 114004, [arXiv:1001.5161 \[gr-qc\]](#).
- [193] E. E. Boos and A. I. Davydychev, “A method for calculating vertex-type Feynman integrals,” *Vestn. Mosk. Univ. (Ser.3)* **28** (1987) 8–12.
- [194] V. A. Fock, *Theorie von Raum, Zeit und Gravitation*. Akademie-Verlag GmbH, Leipziger Straße 3-4, Berlin W1, 1960.
- [195] Y. Itoh, “Equation of motion for relativistic compact binaries with the strong field point particle limit: Third post-Newtonian order,” *Phys. Rev. D* **69** (2004) 064018, [arXiv:gr-qc/0310029](#).
- [196] P. Jaranowski, “Technicalities in the calculation of the 3rd post-Newtonian dynamics,” in *Mathematics of Gravitation, Part II: Gravitational Wave Detection*, A. Królak, ed., pp. 55–63. Banach Center Publications, Vol. 41, Part II, Warszawa, 1997.
- [197] L. Infeld and J. Plebański, *Motion and relativity*. Physical Monographs. Pergamon Press, 4&5 Fitzroy Square, London. W.1., 1960.
- [198] G. Schäfer, “Post-Newtonian methods: Analytic results on the binary problem,” in *Mass and Motion in General Relativity*, L. Blanchet, A. Spallicci, and B. Whiting, eds. Springer, Berlin, 2010. [arXiv:0910.2857 \[gr-qc\]](#).
- [199] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Dover Publications, 1964.
- [200] M. Riesz, “L’intégrale de Riemann-Liouville et le problème de Cauchy,” *Acta Math.* **81** (1949) 1–222.
- [201] M. Riesz, “Erratum: L’intégrale de Riemann-Liouville et le problème de Cauchy,” *Acta Math.* **81** (1949) 223(E).

- [202] A. P. Prudnikov, Y. A. Brychkov, and O. I. Marichev, *More Special Functions*, vol. 3 of *Integrals and Series*. Gordon and Breach, 1986.
- [203] F. Klein, *Vorlesungen über die hypergeometrische Funktion*, vol. 39 of *Grundlehren der mathematischen Wissenschaften*. Springer, 1933.
- [204] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, *NIST Handbook of Mathematical Functions*. Cambridge University Press, Cambridge, 1st ed., 2010.

A Integration Techniques

In this appendix we recapitulate the techniques needed to solve the integrals that appear during our calculation. Most integration techniques are needed for $d = 3$ only, but whenever possible we provide results for generic d here as they may be used for checks. The short-range part of dimensional regularization using an UV-analysis is explained in Sect. 6.4.

A.1 Inverse Laplacians

Inverse Laplacians are necessary to obtain solutions for the fields in the constraints, ϕ and $\tilde{\pi}^i$. As one can see in (43), (45), and (36), the constraint equations always reduce to Poisson-like equations. On the right-hand-side of the equations there is always some source expression appearing. If the source only consists of Dirac deltas or their derivatives the inverse Laplacians can be calculated very easily,

$$\Delta^{-1}\delta_a = -\frac{\Gamma(\frac{d}{2}-1)}{4\pi^{d/2}}\frac{1}{r_a^{d-2}}. \quad (167)$$

Also for some arbitrary power of r_a one can calculate the inverse Laplacian immediately, namely

$$\Delta^{-1}r_a^\alpha = \frac{r_a^{\alpha+2}}{(\alpha+d)(\alpha+2)}. \quad (168)$$

The generalization to the application of multiple inverse Laplacians is straightforward. More interesting is the calculation of inverse Laplacians for powers of r_a and r_b which is a special case of the three-particle integral originally derived in [193, Eq. (7)] and also later given in [69, Eq. above (B14)]. The regular solution of $\Delta^{-1}\frac{1}{r_a r_b}$ in $d = 3$ is well-known and given by

$$\Delta^{-1}\frac{1}{r_a r_b} = \ln s_{ab} = \ln(r_a + r_b + r_{ab}). \quad (169)$$

It was first found in [194], also used in [32] and finally rederived in [69, Eq. (B14)]. (Notice that in [184, Appendix C] a d -dimensional generalization of (169) was given, but its extension to higher inverse Laplacians, i.e. calculation of $\Delta^{-n}(r_a^{2-d}r_b^{2-d})$, is highly non-trivial.) Such solutions are for example necessary for the derivation of $h_{(4)ij}^{\text{TT}}$ [see (117)]. The inverse Laplacians for more general powers of r_a and r_b namely $r_a^m r_b^n$ ($n, m \geq -1$ and n, m odd) in $d = 3$ can be found by using the ansatz

$$\Delta^{-1}r_a^m r_b^n = W_1^{m,n}(r_a, r_b, r_{ab}) + W_2^{m,n}(r_a, r_b, r_{ab}) \ln s_{ab}, \quad (170)$$

where $W_1^{m,n}$ and $W_2^{m,n}$ are polynomials of $(n+m+2)$ -th degree in r_a, r_b , and r_{ab} which consist altogether of $2(\frac{m+n}{2}+2)(m+n+3)$ unknown coefficients. These coefficients have to be fixed by certain consistency conditions

$$\Delta\Delta^{-1} = 1, \quad \Delta_a\Delta^{-1} - \Delta^{-1}\Delta_a = 0, \quad \Delta_b\Delta^{-1} - \Delta^{-1}\Delta_b = 0, \quad (171)$$

where $\Delta_a = \partial_i^{(a)}\partial_i^{(a)}$ denotes the Laplacian with respect to an object coordinate $\hat{\mathbf{z}}_a$. These considerations can be generalized for higher inverse Laplacians. In [27] this technique was extensively used. (Generalizations of the mentioned method are discussed in [195, Sect. V.A.] where the inverse Laplacians are denoted as superpotentials referring to a non-compact source. See also [35, Sects. 5.1., 5.2.] and references therein.)

Before one of the inverse Laplacians discussed above can be applied, one has to get rid of possible $\mathbf{n}_1$ and $\mathbf{n}_2$ vectors. A way to eliminate these $\mathbf{n}$ -vectors is to rewrite them as derivatives with respect to the particle coordinates, which are then commuted with the inverse Laplacian, see e.g. [196] and [78, Appendix C, Eqs. (C6)-(C8)].

A.2 Averaging Procedures

The averaging over the angle variables (necessary for Hadamard's finite part procedure and the local UV-analysis) is given by

$$\int d\Omega_{a,d-1} = \Omega_{a,d-1}, \quad (172a)$$

$$\int d\Omega_{a,d-1} n_a^{i_1} n_a^{i_2} = \frac{1}{d} \Omega_{a,d-1} \delta_{i_1 i_2}, \quad (172b)$$

$$\int d\Omega_{a,d-1} n_a^{i_1} n_a^{i_2} n_a^{i_3} n_a^{i_4} = \frac{1}{d(d+2)} \Omega_{a,d-1} (\delta_{i_1 i_2} \delta_{i_3 i_4} + \delta_{i_1 i_3} \delta_{i_2 i_4} + \delta_{i_1 i_4} \delta_{i_2 i_3}), \quad (172c)$$

$\vdots$

$$\int d\Omega_{a,d-1} n_a^{i_1} \dots n_a^{i_{2k}} = \frac{(d-2)!!}{(d+2(k-1))!!} \Omega_{a,d-1} (\delta_{i_1 i_2} \dots \delta_{i_{2k-1} i_{2k}}), \quad (172d)$$

where (172d) for $d = 3$ coincides with Eq. (A28b) in [100]. $\Omega_{a,d-1}$ is the surface of the d -dimensional unit sphere around point a . We use the $A_{\{i_1 i_2 \dots i_\ell\}}$ notation in the same manner like [100, Appendix A5]. This means $A_{\{i_1 i_2 \dots i_\ell\}} = \sum_{\sigma \in \mathfrak{S}} A_{\sigma(i_1) \dots \sigma(i_\ell)}$ where $\mathfrak{S}$ is the *smallest* set of permutations $(1, \dots, \ell)$ making $A_{\{i_1 i_2 \dots i_\ell\}}$ fully symmetrical in $i_1, \dots, i_\ell$. The factors in front of the deltas come from demanding that the trace over pairwise unit vectors should be one and the integral has the value of the surface of the unit sphere after tracing all unit vectors. An integral over an odd number of unit vectors is zero.

Another necessary averaging procedure is the averaging in a subspace perpendicular to an axis given by a certain vector (this is necessary for reducing integrals of the Riesz-type and the generalized Riesz-type which may have a certain tensor structure to a linear combination of scalar integrals of the appropriate type).

A d -dimensional vector space can be decomposed into a line characterized by a certain vector and a $d - 1$ -dimensional subspace S being perpendicular to the vector. Let this vector be $\mathbf{a}$ with $\mathbf{a}^2 = 1$. Then the averaging of a tensor product of the vector $\mathbf{m}$ lying in the $d - 1$ -dimensional subspace (its components parametrized by angular variables) should give only contributions if the number of vectors is even, namely

$$\langle 1 \rangle_S = 1, \quad (173a)$$

$$\langle m_i \rangle_S = 0, \quad (173b)$$

$$\langle m_i m_j \rangle_S = \alpha \mathfrak{p}_{ij}, \quad (173c)$$

$\vdots$

where $\langle \dots \rangle_S$ denotes the $d - 1$ -dimensional averaging and $\mathfrak{p}_{ij}$ the projector onto the subspace. The projector fulfills the well-known identities

$$\mathfrak{p}_{ij} = \mathfrak{p}_{ji}, \quad (174a)$$

$$a^i \mathfrak{p}_{ij} = 0, \quad (174b)$$

$$\mathfrak{p}_{ij} \mathfrak{p}_{jk} = \mathfrak{p}_{ik}, \quad (174c)$$

and is given by $\mathfrak{p}_{ij} = \delta_{ij} - a^i a^j$. The identity in (173c) is motivated by realizing that $\langle m_i m_j \rangle_S$ fulfills the first two projector identities. This can be rigorously shown under the conditions that $\mathbf{m}$ lies in the subspace perpendicular to $\mathbf{a}$ and $|\mathbf{m}|$ does not depend on the direction of $\mathbf{m}$. If one contracts (173c) with δ_{ij} , then $\mathbf{m}^2$ can be pulled out of the averaging brackets (as it is required to be constant in S) and thus

$$\delta_{ij} \langle m_i m_j \rangle_S = \mathbf{m}^2 = \alpha \mathfrak{p}_{ii}, \quad (175)$$

from which one can conclude that $(\mathfrak{p}_{ii} = d - 1)$

$$\alpha = \frac{\mathbf{m}^2}{d - 1}. \quad (176)$$

One can generalize the above result to

$$\langle m^{i_1} \dots m^{i_{2n}} \rangle_S = \frac{(\mathbf{m}^2)^n (d-3)!!}{(d-3+2n)!!} (\mathbf{p}^{i_1 i_2} \dots \mathbf{p}^{i_{2n-1} i_{2n}}). \quad (177)$$

A.3 Finite Part Integration

In the Hamiltonian density there appear several integrals of the delta-type $\int d^d x f(\mathbf{x}) \delta_a$. The function $f(\mathbf{x})$ is often singular at $\mathbf{x} = \hat{\mathbf{z}}_a$. In $d = 3$ these integrals are often evaluated by the well-known Hadamard's partie-finie (finite part) method (see e.g. [27, 32, 196]). Consider a function $f(\mathbf{x})$ which is well-defined in a neighborhood of $\mathbf{x} = \hat{\mathbf{z}}_a$ and singular at this point. Then it is possible to expand this function in a Laurent series around this point using the auxiliary functions $f_{\mathbf{n}}$,

$$f_{\mathbf{n}}(\varepsilon) := f(\hat{\mathbf{z}}_a + \varepsilon \mathbf{n}) = \sum_{\alpha=-N}^{\infty} a_{\alpha}(\mathbf{n}) \varepsilon^{\alpha}. \quad (178)$$

One defines the zero order coefficient of the Laurent expansion averaged over the $\mathbf{n}$ -vectors as the regularized value of f at $\hat{\mathbf{z}}_a$, namely

$$f_{\text{reg}}(\hat{\mathbf{z}}_a) := \frac{1}{4\pi} \int d\Omega_2 a_0(\mathbf{n}). \quad (179)$$

See (172) for the appropriate averaging formulas. Notice that according to [29, 135] the finite part integration is ambiguous at 3PN because in general

$$(f_1 f_2)_{\text{reg}}(\hat{\mathbf{z}}_a) \neq (f_1)_{\text{reg}}(\hat{\mathbf{z}}_a) (f_2)_{\text{reg}}(\hat{\mathbf{z}}_a), \quad (180)$$

consider e.g. $(\phi_{(2)})^4(\hat{\mathbf{z}}_a)$ vs. $(\phi_{(2)}(\hat{\mathbf{z}}_a))^4$ (see also [197, Appendix 2] for a discussion of this “tweedling of products” property). We will discuss how to avoid this issue in Sect. A.5.

A.4 The Riesz Kernel

Because of the difficulties arising from using Dirac delta distributions as sources of fields in a non-linear theory like General Relativity (e.g. the product of distributions is not well-defined), reformulations of the delta distribution in terms of a function of finite width are also useful, e.g. the Riesz kernel. This kernel tends to a Dirac delta when the regulator (the finite width) tends to zero. The Riesz kernel is given by

$$\delta^{\epsilon_a}(\mathbf{x} - \mathbf{z}_a) = \frac{\Gamma(\frac{d-\epsilon_a}{2})}{\pi^{d/2} 2^{\epsilon_a} \Gamma(\frac{\epsilon_a}{2})} r_a^{\epsilon_a-d}. \quad (181)$$

Thus, in order to get rid of the singularities appearing due to the usage of Dirac deltas, one can replace them by the Riesz kernel and taking the limit $\epsilon_a \rightarrow 0$ for the Riesz kernel regulators after calculating the Hamiltonian. Another advantage of the Riesz kernel is that there will be no distributional contributions to some of the field derivatives mentioned in 6.3. Yet it is important to use Riesz kernels in generic dimension to avoid ambiguous results (as they appear in the finite part method). Unfortunately some of the integrals (in particular inverse Laplacians) for the solutions of the constraints or wave equation have not been solved to date if a Riesz kernel type source is used. Therefore we used the Riesz kernel method only when squares of delta distributions appeared or as a check for the other methods. Another problem of the Riesz kernel is that it breaks the general covariance of the theory explicitly via violating the contracted Bianchi identities $\nabla_{\mu} T^{\mu\nu} = 0$ and one hopes that after the process $\epsilon_a \rightarrow 0$ the covariance is restored. This can be checked by using the Poincaré algebra, see Sect. 8. See [198, above and below Eq. (94)] for a more detailed discussion of this issue.

A.5 Treating Contact Terms

Integration of the delta-type integrands is a little subtle. It was mentioned in [31] that by analysis of Fourier representations of the integrals *within dimensional regularization* one can show that

$$\int d^d x f_1 f_2 f_3 \delta_a = (f_1)_{\text{reg}}(\hat{\mathbf{z}}_a) (f_2)_{\text{reg}}(\hat{\mathbf{z}}_a) (f_3)_{\text{reg}}(\hat{\mathbf{z}}_a), \quad (182)$$

up to and including the formal 3PN order. (But to our knowledge no general proof was given yet.) We evaluated delta-type integrals with the help of this formula and used the d -dimensional Riesz kernel to calculate the regularized values of the functions at the source point appearing on the right hand side. We found that for all cases the regularized values of the fields agreed with results from the usual finite part method in $d = 3$, though this does not hold for products of the fields like $(\phi_{(2)}^4)(\hat{\mathbf{z}}_a)$. Further, the only field leading to nonvanishing finite parts in (93) is $h_{(40)ij}^{\text{TT}}$ and its derivatives.

A.6 Reduction to Scalar Integrals

Besides the delta-type integrals discussed in the previous subsection (or after insertion of Riesz kernels), the only other type of integrals at formal 3PN level is of the form

$$\int d^3 x n_1^{i_1} \dots n_1^{i_k} \cdot n_2^{j_1} \dots n_2^{j_\ell} \cdot r_1^\alpha r_2^\beta s_{12}^\gamma. \quad (183)$$

The s_{12} is introduced from derivatives of $\ln s_{12}$, which in turn arise from certain inverse Laplacians. This subsection provides a formalism to reduce these integrands with complicated tensor structure to a linear combination of pure scalar integrands of the generalized Riesz-type (discussed in the next subsection). It is only possible to rewrite the vectors as derivatives with respect to the particle coordinates applied to a function of the type $r_1^\alpha r_2^\beta s_{12}^\gamma$ (as for the inverse Laplacians) for the case $\gamma = 0, 1, 2, \dots$, because the s_{12} also depends on the particle coordinate. For integer $\gamma = n \geq 0$

$$s_{12}^n = (r_1 + r_2 + r_{12})^n = \sum_{k=0}^n \sum_{\ell=0}^k \binom{n}{k} \binom{k}{\ell} r_1^\ell r_2^{k-\ell} r_{12}^{n-k}, \quad (184)$$

and therefore the mentioned functions can be reduced to a linear combination of products of the form $r_1^\alpha r_2^\beta r_{12}^\gamma$. This product structure is crucial to rewrite the vectors in (183) into particle derivatives. Thus one has to use another method to eliminate the vectors from the integrand for $\gamma < 0$ where the functions cannot be reduced to a product form. In [27] a method is mentioned which can be used to get rid of angle integrations resulting from the vectors by using an averaging procedure of the integrand in prolate spheroidal coordinates. We will present this procedure here in a slightly different way.

First of all one has to get rid of, e.g., the $\mathbf{n}_2$ vectors by using the identity

$$\mathbf{n}_2 = \frac{r_1}{r_2} \mathbf{n}_1 + \frac{r_{12}}{r_2} \mathbf{n}_{12}. \quad (185)$$

Afterwards one still has to eliminate $\mathbf{n}_1$ vectors from the integrand. By using $\mathbf{n}_1 = (\mathbf{x} - \hat{\mathbf{z}}_1)/r_1$ and the orthogonal decomposition of $\mathbf{n}_1$ with respect to $\mathbf{n}_{12}$ one gets

$$\mathbf{n}_1 = (\mathbf{n}_1 \mathbf{n}_{12}) \mathbf{n}_{12} + \mathbf{n}_1^\perp, \quad n_1^{\perp i} = \mathfrak{P}^{ij} n_1^j, \quad (186)$$

where

$$(\mathbf{n}_1 \mathbf{n}_{12}) =: A = \frac{1}{2r_1 r_{12}} (r_2^2 - r_1^2 - r_{12}^2), \quad (187)$$

$$|\mathbf{n}_1^\perp| =: B = \frac{1}{2r_1 r_{12}} \sqrt{[(r_1 + r_2)^2 - r_{12}^2] [r_{12}^2 - (r_1 - r_2)^2]}. \quad (188)$$

and $\mathfrak{P}^{ij} = \delta_{ij} - n_{12}^i n_{12}^j$ is the projector onto the subspace perpendicular to $\mathbf{n}_{12}$. Equation (187) can be derived by considering $r_2^2 = |\mathbf{x} - \hat{\mathbf{z}}_2|^2 = |\mathbf{x} - \hat{\mathbf{z}}_1 + \hat{\mathbf{z}}_1 - \hat{\mathbf{z}}_2|^2$ and expressing it in terms of r_1 , r_{12} and A

itself. B is given by $B^2 = n_1^i \mathfrak{P}^{ij} n_1^j = 1 - A^2$ where we used the projector condition, $n_{12}^i \mathfrak{P}^{ij} = 0$, (186), and (187). Now a tensor product of such vectors can also be decomposed via (186) as

$$\int d^d x T_{i_1 \dots i_k} n_1^{i_1} \dots n_1^{i_k} = \int d^d x T_{i_1 \dots i_k} (n_1^\perp + A n_{12})^{i_1} \dots (n_1^\perp + A n_{12})^{i_k}, \quad (189)$$

where $T_{i_1 \dots i_k}$ collects all constant tensors like linear momenta, spin tensors, and $\mathbf{n}_{12}$ vectors. Notice that neither A nor B depend on the direction of $\mathbf{n}_1^\perp$. Furthermore the integration extends over the whole $\mathbb{R}^d$ such that one can rewrite the integrals of the type

$$\int d^d x S_{i_1 \dots i_k}(r_1, r_2, r_{12}) n_1^{\perp i_1} \dots n_1^{\perp i_k}, \quad (190)$$

in a more convenient way after a coordinate transform to prolate spheroidal coordinates. The Cartesian coordinates in terms of the prolate spheroidal coordinates $(\xi, \eta, \phi_1, \dots, \phi_{d-2})$ are given by (generalization of the coordinates given in [199, pp. 752])

$$x_1 = f \xi \eta, \quad (191a)$$

$$x_2 = f \sqrt{(\xi^2 - 1)(1 - \eta^2)} \cos \phi_1, \quad (191b)$$

$$x_3 = f \sqrt{(\xi^2 - 1)(1 - \eta^2)} \sin \phi_1 \cos \phi_2, \quad (191c)$$

$\vdots$

$$x_{d-1} = f \sqrt{(\xi^2 - 1)(1 - \eta^2)} \sin \phi_1 \dots \cos \phi_{d-2} \quad (191d)$$

$$x_d = f \sqrt{(\xi^2 - 1)(1 - \eta^2)} \sin \phi_1 \dots \sin \phi_{d-2} \quad (191e)$$

where $-1 < \eta < 1$; $1 < \xi < \infty$; $0 \leq \phi_{d-2} \leq 2\pi$; $0 \leq \phi_1, \dots, \phi_{d-3} \leq \pi$. ξ and η can be related to distance variables r_1, r_2, r_{12} by using

$$\xi = \frac{r_1 + r_2}{r_{12}}, \quad \eta = \frac{r_1 - r_2}{r_{12}}, \quad f = \frac{r_{12}}{2}. \quad (192)$$

Now one can easily perform the transformation of the volume element which is given by

$$d^d x = f^d (\xi^2 - \eta^2) d\xi d\eta d\Omega_{d-2}. \quad (193)$$

Thus we can perform an averaging over the $d - 1$ -dimensional subspace perpendicular to $\mathbf{n}_{12}$ by transforming the volume form

$$\begin{aligned} \int d^d x S_{i_1 \dots i_k}(r_1, r_2, r_{12}) n_1^{\perp i_1} \dots n_1^{\perp i_k} &= \int d\xi d\eta f^d (\xi^2 - \eta^2) S_{i_1 \dots i_k}(\xi, \eta) \\ &\times \int d\Omega_{d-2} n_1^{\perp i_1} \dots n_1^{\perp i_k}, \\ &= \int d\xi d\eta \Omega_{d-2} f^d (\xi^2 - \eta^2) S_{i_1 \dots i_k}(\xi, \eta) \langle n_1^{\perp i_1} \dots n_1^{\perp i_k} \rangle, \\ &= \int d^d x S_{i_1 \dots i_k}(r_1, r_2, r_{12}) \langle n_1^{\perp i_1} \dots n_1^{\perp i_k} \rangle, \end{aligned} \quad (194)$$

where $\langle n_1^{\perp i} \rangle = 0$ and $\langle n_1^{\perp i} n_1^{\perp j} \rangle = \frac{B^2}{d-1} \mathfrak{P}^{ij}$. Notice that $\mathfrak{P}^{ij}$ has the same properties like $\mathfrak{p}_{ij}$ and $\langle \dots \rangle$ the same like $\langle \dots \rangle_S$ in A.2. Further notice that the average in the integrand only depends on r_1, r_2 , and r_{12} . Thus one is able to reduce all integrals of the generalized Riesz-type containing $\mathbf{n}$ -vectors to a linear combination of scalar integrals. Notice that in the case $\gamma \geq 0$ the method from the present subsection is also advantageous over rewriting $\mathbf{n}$ -vectors as derivatives. The latter can lead to new singularities, logarithms, or even a need for a new kind of regulator, e.g., for an expression like

$$\int d^3 x n_a^i n_a^j n_a^k r_a^{-3}. \quad (195)$$

By using the method of the present subsection these issues are avoided.

A.7 Integration using Generalized Riesz-Formula

After reduction to scalar integrals one is left with integrals of the form $\int d^3x r_1^\alpha r_2^\beta s_{12}^\gamma$. For integer $\gamma \geq 0$ one can use (184) and end up with integrals whose solution were found by M. Riesz for arbitrary d [200, 201], namely

$$\int d^d x r_1^\alpha r_2^\beta = \pi^{d/2} \frac{\Gamma(\frac{\alpha+d}{2})\Gamma(\frac{\beta+d}{2})\Gamma(-\frac{\alpha+\beta+d}{2})}{\Gamma(-\frac{\alpha}{2})\Gamma(-\frac{\beta}{2})\Gamma(\frac{\alpha+\beta+2d}{2})} r_{12}^{\alpha+\beta+d}. \quad (196)$$

For $d = 3$ a generalization of the Riesz-formula was found by P. Jaranowski and G. Schäfer in [27],

$$\begin{aligned} \int d^3x r_1^\alpha r_2^\beta s_{12}^\gamma &= 2\pi \frac{\Gamma(\alpha+2)\Gamma(\beta+2)\Gamma(-\alpha-\beta-\gamma-4)}{\Gamma(-\gamma)} \left(I_{1/2}(\alpha+2, -\alpha-\gamma-2) \right. \\ &\quad \left. + I_{1/2}(\beta+2, -\beta-\gamma-2) \right. \\ &\quad \left. - I_{1/2}(\alpha+\beta+4, -\alpha-\beta-\gamma-4) - 1 \right) r_{12}^{\alpha+\beta+\gamma+3}, \end{aligned} \quad (197)$$

with $I_{1/2}(x, y) := \frac{B_{1/2}(x, y)}{B(x, y)}$ the incomplete regularized Euler Beta function. One can express the incomplete Beta function (Euler integral of the first kind) $B_{1/2}(x, y)$ in terms of the Gauß hypergeometric function ${}_2F_1$

$$B_{1/2}(x, y) = \frac{1}{2x} {}_2F_1\left(1-y, x; x+1; \frac{1}{2}\right) = \frac{1}{x} {}_2F_1(x+y, x; 1+x; -1). \quad (198)$$

The function $B(x, y)$ represents the Euler Beta function or Euler integral of the second kind with $B(x, y) = \frac{\Gamma(x+y)}{\Gamma(x)\Gamma(y)}$. It turns out that the regularization procedure mentioned in [27, Eq. (B21), (B23), and (B24)] only modifies α and β to be non-integer via the introduced analytical regulators $\mu\epsilon$ and $\nu\epsilon$. So one can in principle simplify the formula given above by the assumption that $\gamma \in \mathbb{Z}$, and α and β being arbitrary. Positive integer powers of s_{12} can be handled through (184) and (196). Thus the only relevant powers of s_{12} are the negative integer ones.

Equation (198) leads to hypergeometric functions of the type ${}_2F_1(-\gamma, z; z+1; -1)$ in (197), where z depends on α and β , namely

$$\begin{aligned} \int d^3x r_1^\alpha r_2^\beta s_{12}^\gamma &= 2\pi\Gamma(2+\alpha)\Gamma(2+\beta)\Gamma(-4-\alpha-\beta-\gamma) \left\{ -\frac{1}{\Gamma(-\gamma)} \right. \\ &\quad \left. + \frac{{}_2F_1(-\gamma, 2+\alpha; 3+\alpha; -1)}{\Gamma(3+\alpha)\Gamma(-2-\alpha-\gamma)} + \frac{{}_2F_1(-\gamma, 2+\beta; 3+\beta; -1)}{\Gamma(3+\beta)\Gamma(-2-\beta-\gamma)} \right. \\ &\quad \left. - \frac{{}_2F_1(-\gamma, 4+\alpha+\beta; 5+\alpha+\beta; -1)}{\Gamma(5+\alpha+\beta)\Gamma(-4-\alpha-\beta-\gamma)} \right\} r_{12}^{\alpha+\beta+\gamma+3}. \end{aligned} \quad (199)$$

So for negative integer gamma $\gamma = -n$ one can express the solution of the integral in terms of ${}_2F_1(n, z; z+1; -1)$. It is well-known that [202]

$${}_2F_1(0, z; z+1; -1) = 1, \quad (200)$$

$${}_2F_1(1, z; z+1; -1) = \frac{z}{2} \left[\psi\left(\frac{z+1}{2}\right) - \psi\left(\frac{z}{2}\right) \right], \quad (201)$$

where ψ is the Digamma function. From these two formulas and the contiguous relation of the Gauß hypergeometric function [199, 203, 204]

$$\begin{aligned} 0 &= (c-a){}_2F_1(a-1, b; c; z) + (2a-c+(b-a)z){}_2F_1(a, b; c; z) \\ &\quad + a(z-1){}_2F_1(a+1, b; c; z), \end{aligned} \quad (202)$$

one obtains the recursive relation

$${}_2F_1(n, z; z+1; -1) = \frac{1}{2(n-1)} \left[(z - (n-2)) {}_2F_1(n-2, z; z+1; -1) + (3n - 2(2+z)) {}_2F_1(n-1, z; z+1; -1) \right]. \quad (203)$$

From these relations one can show per induction the general structure of the hypergeometric function to be

$${}_2F_1(n, z; z+1; -1) = \sum_{k=0}^{n-1} a_k^{(n)} z^k + \frac{(-1)^{n-1}}{2(n-1)!} \left[\psi\left(\frac{z+1}{2}\right) - \psi\left(\frac{z}{2}\right) \right] \prod_{k=0}^{n-1} (z-k). \quad (204)$$

Unfortunately the coefficients $a_k^{(n)}$ can only be obtained by complicated recursive relations between different k and n following from (203) and we can only give an explicit form for some of them

$$a_0^{(n)} = 0, n \geq 2 \quad \text{and} \quad a_0^{(0)} = 1, a_0^{(1)} = 0, \quad (205)$$

$$a_{n-1}^{(n)} = \frac{(-1)^n}{2(n-1)!}, n \geq 2, \quad (206)$$

$$a_{n-2}^{(n)} = \frac{(-1)^n (1+n-n^2)}{4(n-1)!}, n \geq 3. \quad (207)$$

Nevertheless the mentioned recursion relations can be used to cache all Gauß hypergeometric functions. MATHEMATICA is able to handle limits, series expansions and derivatives of the arising Digamma functions very well and thus all occurring integrals in the binary Hamiltonian to linear order in the spin variables can be solved.